procedure was CT scan, of which 132 (32%) were potentially redundant, for a rate of 7.4% (5.0-10.6%) per 100,000 member months. Rates will be calculated for a similar period following the HIE implementation and compared to baseline rates to assess the impact. **Discussion:** The HIE represents a new tool that can potentially reduce the rates of redundant testing in the ED setting. The process of defining redundant tests is a critical component of this kind of analysis. More work is needed in this area to define, standardize, and validate the "rules" for categorizing tests as redundant.

Keywords: Health Information Exchange; Test/Study Redundancy; Health Informatics

doi:10.3121/cmr.2012.1100.ps1-50

PS1-51:

Effect of Integrated Health Information Exchange on the Duplication of Radiological Services

Christine Bredfeldt¹

¹Kaiser Permanente Mid-Atlantic, MAPMG

Background/Aims: Unnecessary duplication of diagnostic procedures such as x-rays increases healthcare costs and jeopardizes patient safety. Electronic health records (EHRs) have the potential to reduce unnecessary duplication of diagnostic studies by improving timely access to the original test results. However, when follow-up care occurs in a different institutional setting, providers may not have access to the same EHR. In such cases, electronic health information exchange (HIE) could improve the quality and timeliness of information available to the follow-up provider, thereby reducing the need to order duplicative diagnostic tests. Our primary goal in this work was to determine if members with diagnoses of bone fractures were less likely to receive follow-up x-rays when findings from the initial diagnostic imaging study were available in the electronic health record. Our secondary goal was to estimate the economic costs and patient safety repercussions of duplicate imaging studies. **Methods:** This retrospective cohort study of Kaiser Permanent members in the Mid-Atlantic States (KPMAS) compared the rate of duplicate x-rays in patients with a diagnosis of bone fractures from the Emergency Department (ED) or from outpatient care between 2006 and 2010. We used the Medicare fee schedule to estimate costs for the total episode of care. We estimated radiation exposure by identifying all relevant radiology procedures during the two months following the index event and assigning radiation levels based on published estimates for each procedure.

Results: The study included approximately 50,000 patients from KPMAS with a bone fractures diagnosis. 20% of patients were initially seen in the ED, while 80% of patients were seen in outpatient care. Patients initially seen at a KPMAS outpatient facility were more likely than patients initially seen in the ED to have x-ray findings recorded in the KPMAS EHR prior to their follow-up visit. These patients were also significantly less likely to have duplicated (same or similar) radiology procedures ordered at follow-up visits. **Discussion:** When the findings from preliminary imaging studies are available in the EHR, providers are less likely to order similar x-rays at follow-up visits. This reduction in duplicate radiology procedures significantly reduces costs and increases patient safety.

Keywords: Electronic Health Records; Health Information Exchange; Health Informatics

doi:10.3121/cmr.2012.1100.ps1-51

PS1-52:

Charting the Course: Adjudicating Fractures in Network Enrollees Using the Electronic Medical Record

Jane Grafton¹; Susan Ott²; Julie Endres¹; Christel Kratochvil¹; Rebecca Hubbard¹; Laura Ichikawa¹; Jacqueline St. John¹; Belinda Operskalski¹; Leslie Spangler¹; Julia Anderson¹; Eileen Rillamas-Sun¹; Andrea LaCroix³; Delia Scholes¹

¹Group Health Cooperative; ²University of Washington; ³Fred Hutchinson Cancer Research Center

Background/Aims: In an era of increasing openness in health plans, use of electronic medical records (EMR) for detailed chart abstraction for all enrollees poses challenges. At Group Health Cooperative (GH), EMR data

are most complete for enrollees in the Group Practice Division (GPD). However, all GH enrollees (from both the GPD and the Contracted Network Division (CND) have EMRs. CND providers do not have access to patients' GH EMR. Information for CND enrollees is only entered when the patient or provider interacts with the GPD, such as a patient receiving GPD specialty care while receiving CND primary care. In the interest of achieving a larger sample size for an ongoing GH-based case-control study (FOCUS), we evaluated the feasibility of including CND enrollees by determining the proportion with sufficient EMR information for fracture adjudication. **Methods:** The FOCUS study is adjudicating and recruiting incident osteoporotic fracture cases in women across the menopausal transition using the GPD. We used identical automated sampling methods to identify, via ICD-9 codes, a pool of potential adjudicable fracture cases among CND enrollees. **Results:** For the GPD, from a denominator of 58,328 age-eligible women, 2,310 (4%) were identified with fractures January 2008-March 2011. For the CND, among 38,041 age-eligible women, 1,040 (3%) were identified with fractures. Following automated exclusions, 944 CND potential cases remained. Initial review of these 944 EMRs found that 685 charts had no data; 54 had data unrelated to fracture; 145 had sufficient fracture-related data to adjudicate; and 60 had some fracture-related data. Preliminary chart adjudication of these last two groups found that 88% (127/145) of the "complete" EMR group and 10% (6/60) of the "partially complete" group had qualifying fractures. **Discussion:** About 23% of GH network enrollees with automated fracture codes had evaluable EMR data. Preliminary adjudication has identified 65% (133/205) as additional potential FOCUS cases to contact for interviews. It appears that, even for a study requiring sufficient chart information to adjudicate a health outcome, the CND can contribute cases for study inclusion. We also have developed network control selection methods. Still to be ascertained at the survey step are continued eligibility and study participation.

Keywords: Electronic Medical Record; Fracture Adjudication; Health Informatics

doi:10.3121/cmr.2012.1100.ps1-52

CB4-02:

Validation of a Natural Language Processing Protocol for Detecting Heart Failure Signs and Symptoms in Electronic Health Record Text Notes

Roy Byrd¹; Steven Steinhilb²; Walter Stewart²; Zahra Daar²; Shahram Ebadollahi¹; Jimeng Sun¹

¹IBM; ²Geisinger Health System

Background/Aims: Early detection of Heart Failure (HF) could help change the natural history of one of the most costly and disabling conditions among older patients through early intervention with treatments. We describe natural language processing (NLP) methods and validation results for detection of major and minor Framingham criteria in electronic health record (EHR) clinical notes as a first step towards early detection of HF. **Methods:** Data for this study were from 6,355 primary care incident heart failure cases identified using Geisinger EHR data between 2001 and 2010 and from 26,052 controls. Text data were from Office Visit, Case Manager, and Radiology notes in the four-year period before diagnosis of HF in cases and a comparable period for controls. Text analysis comprised: 1) application-independent analytics for paragraph and sentence identification, tokenization, dictionary look-up, morphological analysis, and part-of-speech tagging; 2) dictionaries and grammars for 9 major and 6 minor Framingham criteria, and for the compound noun phrases, negated contexts, and counterfactual contexts in which criteria are mentioned; and 3) text analysis engines to disambiguate criteria mentions, to apply constraint rules (e.g., word co-occurrence), and to decide whether identified signs and symptoms are being affirmed or denied. During training, NLP results were iteratively validated against the judgments of a cardiologist. The final system was tested against gold standard annotations reflecting the consensus judgments of four trained coding experts. **Results:** Over an 8-week period of iterative training, the NLP results on a dataset of 65 encounter notes improved from a positive predictive value (PPV) of 0.741 and a sensitivity (Se) of 0.504 to a final PPV of 0.931 and a Se of 0.939. On the consensus gold standard dataset of 200 cases and 200 controls, the fully-trained system achieved a PPV of 0.925 and a Se of 0.896. **Discussion:** The NLP system can accurately identify affirmations and denials of Framingham criteria, which can directly reduce physicians workload for