PS1-56:

Free Generic Medication Programs and the Patients that Take Advantage of Them: A Look at Metformin

Melissa Butler¹; Anping Chang¹; Jennifer Kuntz²

¹Kaiser Permanente Southeast; ²Kaiser Permanente Northwest

Background/Aims: In 2006, pharmacy retailers began offering low cost generic medications at costs ranging from \$4 for a 30-day supply to \$10 for a 90-day supply. One is now offering free generic metformin. There are considerable cost savings for patients who take advantage of these programs. However, filling medications at multiple pharmacies could lead to incomplete medication histories within pharmacy and insurance records thus making it difficult identify drug interactions and to address quality initiatives. This study will measure trends in requests for prescriptions to be filled within an integrated health system pharmacy versus an outside pharmacy retailer before and after the availability of free metformin in March 2010. Characteristics of patients who switched from prescriptions for internal fills to external fills after free metformin was available will also be examined. **Methods:** Electronic medication prescription order data from January 2009 to December 2011 will be used in the study. Patients with diabetes will be identified by two outpatient visits, one emergency department visit, or one hospitalization with a diagnosis code for diabetes and use of an oral antidiabetes medication. Monthly metformin prescription rates will be compared to generic sulfonylurea prescription rates; sulfonylureas are not available for free. Patients will be included in the analysis if they have 12 months of continuous enrollment with no evidence of either a fill or a prescription order for metformin or a sulfonylurea during that time. An interrupted time series analysis with segmented regression will be used to study the impact of free generic metformin on requests for outside prescription fills. A logistic regression difference-in deference model will be used to study the impact of the availability of free metformin on pharmacy switching. We will also examine the relevance of the following characteristics on switching, demographics, health plan product, number of medications, the day supply usually filled, percent of medications that are generic, copay, and distance to medical office building from residence. **Results:** Analysis is ongoing. **Discussion:** This study will help measure the extent of external medication fills and characteristics of patients who fill medications externally in a market where medications are available for free.

Keywords: Generic Medication; Medication Costs; Health Services Research

doi:10.3121/cmr.2012.1100.ps1-56

PS1-58:

Clinicians' Use of Automated Reports of Estimated Glomerular Filtration Rate: A Qualitative Study

David Smith¹; Jennifer Schneider¹; Micah Thorp¹; Suma Vupputuri²; Jessica Weiss^{1,3}; Eric Johnson¹; Adrienne Feldstein¹; Amanda Petrik¹; Xiuhai Yang¹; Susan Snyder⁴

¹Kaiser Permanente Northwest; ²Kaiser Permanente Southeast; ³Oregon Health Sciences University; ⁴Centers for Disease Control and Prevention

Background/Aims: Improving both awareness and identification of chronic kidney disease (CKD) is important among primary care providers so they can provide appropriate clinical care; one method that has been widely embraced is the use of automated reporting of estimated glomerular filtration rate (eGFR) by clinical laboratories. We undertook a qualitative study to examine how clinicians responded to the onset of automated reporting of eGFR at Kaiser Permanente Northwest (KPNW). **Methods:** We conducted 19 semi-structured interviews among primary care clinicians including both physicians (n=13) and allied health providers (n=6), recruited from KPNW, a non-profit health maintenance organization. Interviews were approximately 30-45 minutes in length, audio-recorded and transcribed. A standard, qualitative content analysis approach was employed. **Results:** Clinicians generally held favorable views toward automated eGFR reporting as useful information they wish they had easy access to previously. Most (n=15) reported using the eGFR in conjunction with serum creatinine in their clinical decision-making. Benefits of the automated eGFR reporting included saving valuable clinic time in not having to calculate it on their own, increased awareness of and

attention to tracking CKD stages, and improved patient care. Increased patient education and counseling on CKD prevention and kidney health was reported by 10 of the clinicians interviewed, and 14 reported minor increases in their referral to Nephrology. Challenges experienced as a result of the automated eGFR reporting included managing patient confusion or fear from the reporting, and increased workload. Suggested improvements included offering yearly trainings on CKD management, providing regular feedback from Nephrology about appropriateness of clinician referral request, improving the integration of eGFR value with other tools in the electronic medical record, and providing tools to facilitate discussion of eGFR findings with patients. **Discussion:** Overall, clinicians used automated eGFR as a tool to help identify CKD stage, educate and counsel patients about their kidney function and health, and make treatment decisions. The manner in which clinicians use eGFRs appears to be more complex than previously understood, and our study illustrates some of the efforts that might be usefully undertaken (e.g. specific clinician education) when encouraging further promulgation of eGFR reporting and usage.

Keywords: Chronic Kidney Disease; Glomerular Filtration Rate; Health Services Research

doi:10.3121/cmr.2012.1100.ps1-58

PS1-59:

Collecting Biomedical Specimens in Health Research

Kevin Ulrich¹

¹Group Health, Research Institute

Survey research organizations are increasingly being tasked with obtaining consent and, in some cases, collecting biological specimens from potential respondents. In order to obtain high response rates in these types of studies, it is crucial that survey methodologists investigate best practices to improve response in the collection of these types of data. Under what circumstances are participants willing to consent to these procedures? What methods can be utilized to best facilitate the collection of these types of data? This poster reviews outcomes from multiple studies conducted by the Survey Research Program at Group Health Research Institute, which required the collection of various biological specimens. Recruitment procedures, methodologies for obtaining consent, response rates, characteristics of participants and non-participants, and recommendations for best practices are discussed.

Keywords: Biomedical Specimens; Response Rates; Health Services Research

doi:10.3121/cmr.2012.1100.ps1-59

PS1-61:

Implementing CHES eHealth Breast Cancer Support in Population-based Care

Alanna Kulchak Rahm¹; James Dearing¹; Breanne Barela¹; Suzanne Pingree²; Robert Hawkins²

¹Kaiser Permanente Colorado; ²University of Wisconsin Madison

Background/Aims: The woman who is newly diagnosed with breast cancer suddenly faces significant new needs. She must learn about her disease and its treatment, cope with emotional distress and interpersonal issues, and make decisions about her treatment, work and family obligations. To answer the need for information and support, an Interactive Cancer Communication Systems (ICCS) called CHES has been well-researched and validated to provide these women access to integrated information for decision-making, behavior change, and emotional support. **Methods:** This newly-funded dissemination and implementation project will offer CHES to all newly diagnosed breast cancer patients as part of standard care in two Denver healthcare systems, Kaiser Permanente and Exempla St. Joseph's Hospital. This project is designed to evaluate the real-world implementation of this new technology by answering the "how much" and "why" questions of utilization at both the institutional and patient levels. We will assess how effectively CHES is offered to patients at each institution as part of standard care, what proportion of patients utilize the CHES system and how often, why some women use it and some do not, to what extent women use the system; how staff perceptions and actions affect patient perceptions and use, if CHES use is related to patient satisfaction and quality of life, and how