for a cohort of Kaiser Permanente Northwest (KPNW) members receiving surgical procedures of the cervix. **Methods:** The HMORN Virtual Data Warehouse (VDW) was used to obtain health plan enrollment, diagnosis, procedure, and demographic data for a cohort of 14- to 53-year-old KPNW women during the study years, 1998-2009. Unexposed women were matched to exposed women on age and calendar year. We performed logistic regression to model the propensity of receiving surgical procedures of the cervix. Potential covariates included smoking status, race and ethnicity, sexually transmitted infections (STI), immunocompromised status, length of health plan enrollment, and number of encounters per enrollment year. Propensity score balance was assessed using histograms and boxplots. The common support condition was imposed. Matching on propensity scores was performed using a greedy 5 to 1 digit match. **Results:** There were 86,898 women in our cohort. Women receiving a surgical procedure were more likely to be current tobacco users, immunocompromised, have a history of STI, have longer prior enrolment in the health plan, and have more healthcare encounters per enrolment year than unexposed women. In our cohort, propensity scores for receiving invasive cervical procedures for exposed women ranged from 0.02 to 0.56 (n=4138; mean=0.08; median=0.06), whereas propensity scores for unexposed women ranged from 0.01 to 0.99 (n=82760; mean=0.05; median=0.04). We excluded 1,819 (2%) unexposed women based on the common support condition. A 1:3 match ratio produced the best matched population. No matches were found for 43 (1%) of our exposed population. **Discussion:** Use of the HMORN VDW enabled us to create propensity scores for estimating the probability of receiving cervical procedures. Matching and imposing the common support condition decreased our study population minimally. However, women excluded from analysis may be intrinsically different from those who remained. One limitation to our study is that some potential variables we wished to include in our model, such as marital status, were not available in HMORN VDW data which may introduce bias.

Keywords: Propensity Score; Probability; Mega-epidemiology and Methods

doi:10.3121/cmr.2012.1100.ca5-02

CA5-03:

Bringing Data Closer to Investigators

Nicolas Beaulieu¹; Tiffany Woodworth¹; James Marshall¹; Ashley Wong¹; Lisa Trebino¹; Jeffrey Brown¹

¹Harvard Pilgrim Health Care Institute

Background/Aims: To allow rapid response to FDA data inquiries, Mini-Sentinel Operations Center (MSOC) has developed seven modular programs (MP) for use within FDA's Mini-Sentinel Pilot, a large, multi-institutional distributed database with 17 data partners (DP) designed to support the agency's efforts to develop an active surveillance system for monitoring medical product safety in the United States. **Methods:** MPs are SAS programs designed to run against the Mini-Sentinel Common Data Model executing a set of common instructions for routine questions. MPs address questions commonly asked within the context of medical product safety assessments such as the number of prevalent and incident users of a medical product, the number of dispensings and days supply of specific drug products, the number of patients with a specific diagnosis or procedure, and the number of outcomes identified during exposure person-time. MP output includes standardized summary-level counts and are presented overall and stratified by user-defined age group, sex and calendar year. A set of input forms allow investigators to easily define parameters (e.g., lengths of drug washout period and look back period for pre-existing conditions, minimum treatment episode duration, allowable gap between dispensings). DPs are expected to respond to MP requests within five business days. Results are returned to MSOC via a secure portal and subsequently a summarized report is generated and returned to FDA requesters. **Results:** Between February and October 2011, MSOC has completed 15 MP queries in response to specific FDA questions for a total of 235 sets of results returned by participating DPs of which 78.3% were completed within the allotted 5 days. The mean response time across all queries was 3.8 days. Average turnaround time for MSOC to create a summarized report for FDA was 7.0 business days. **Discussion:** A key benefit of MP-based data request system is that simple summarized data are easily available to investigators. The process will be assessed and modified, as use of the system generates lessons and new

approaches to improve efficiency. Planned future developments include addition of new data elements to MPs (e.g., lab values) and decision tree tool to match investigators' question to the appropriate MP.

Keywords: Distributed database; Modular Program; Mega-epidemiology and Methods

doi:10.3121/cmr.2012.1100.ca5-03

CA5-04:

Identifying Pregnancy Episodes in the Vaccine Safety Datalink

Allison Naleway¹; Rachel Gold¹; Michelle Henninger¹; Samantha Kurosky¹; Karen Riedlinger¹; James Nordin²; Elyse Kharbanda²; Eric Weintraub³

¹Kaiser Permanente; ²HealthPartners; ³Centers for Disease Control and Prevention

Background/Aims: The need for research on the safety of vaccination during pregnancy is widely recognized. Large, population-based data systems like the Vaccine Safety Datalink (VSD) may be useful for this research, but identifying pregnancies using electronic medical record (EMR) and claims data can be challenging. **Methods:** We modified an existing pregnancy identification algorithm originally developed by Kaiser Permanente Northwest to identify pregnancy outcomes and dates using the standardized VSD data files. We validated the algorithm by calculating the percent agreement in pregnancy outcome type, end date, and gestational age between the algorithm and manual medical record review. At each site, we randomly sampled 15 episodes within four outcome type strata (live births, spontaneous abortions, elective abortions, and other pregnancy outcomes) for a total of 60 episodes per site. Seven of eight VSD sites participated. **Results:** We identified 595,929 pregnancy episodes ending in 2002-2006 among women 12-55 years of age. Of these pregnancies, 75% ended in live births, 12% in spontaneous abortions, and 9% in elective abortions. We were able to confirm a pregnancy on or near the algorithm-specified pregnancy start and end dates for 99% of live births, 93% of spontaneous abortions, 92% of elective abortions, and 90% of other outcomes. The agreement between the algorithm-identified and the abstractor-identified outcome date ranged from 70% (elective abortion) to 96% (live birth) depending on outcome type. When gestational age was available in the EMR, agreement ranged from 82% (other) to 98% (live birth) depending on outcome type. **Discussion:** The VSD algorithm accurately identifies pregnancy episodes across participating sites using the standardized VSD data files. Additional manual record review may be needed to improve the precision of the pregnancy date estimates depending on specific study needs. This algorithm will allow us to conduct large, population-based studies of the safety of vaccination during pregnancy.

Keywords: Pregnancy; Vaccine Safety; Mega-epidemiology and Methods

doi:10.3121/cmr.2012.1100.ca5-04

CA5-05:

A Visual Method to Rapidly Facilitate Case Identification Among a Medically Complex Cohort

Mary Kay Theis¹; G. Craig Wood²; Emily Westbrook¹; Glenn Gerhard²; David Arterburn¹

¹Group Health Cooperative; ²Geisinger Health System

Background/Aims: Comparative effectiveness research studies often require researchers to make inference about the relationship between exposure to various treatments over time and health outcomes. In these studies it can be a challenge to isolate the effect of the exposure from other concurrent treatments and health conditions. For example, in the study of the effect of different antipsychotic medications on body weight, concurrent exposures of oral steroids, insulin, pregnancy, and cancer diagnosis can have important confounding effects. The temporal relationships between these exposures may be difficult to assess in large samples; therefore we sought a way to graphically display exposures and outcomes in a single figure to facilitate rapid screening of potential cases. **Methods:** Researchers at Geisinger and Group Health identified all adults who a) received one or more prescription for a 2nd/3rd generation antipsychotic medication from 2004-2009 and b) had two or more weight measures in close proximity to those treatments. For each adult who had gained >15% body weight after initiating drug treatment