Accepted By Abstract Software. **Discussion:** The proportion of children and adolescents with both normal BP and normal BMI was 46.7% and decreased with age, while the proportion with both elevated BP and BMI rose from 7.8% to 20.0% with increasing age. Clinical and public policy initiatives to reduce the very high frequency of these risk factors in U.S. children and adolescents should be carefully considered.

Keywords: Blood Pressure; BMI; Prevention and Wellness

doi:10.3121/cmr.2012.1100.cb8-01

CB8-02:

Screening Dual Energy X-Ray Absorptiometry (DXA): Are We Neglecting Women at High Risk of Hip Fracture?

Wendolyn Gozansky¹; Christina Clarke¹; Deanna McQuillan¹; Jennifer Ellis¹

¹Kaiser Permanente Colorado

Background/Aims: Screening DXA is recommended for women aged 65+ to identify individuals likely to benefit from treatments to prevent osteoporotic fracture. This strategy is most cost-saving in women 75+, but DXA use remains low among older women. We determined whether estimates of absolute fracture or fall risk impact DXA screening. **Methods:** Using administrative data, we performed a cross-sectional analysis of women aged 65+ with ≥ 2 years of enrollment in Kaiser Permanente Colorado during 2005-2009. DXA was determined by procedure code or scan documentation prior to the end of observation. To isolate screening, we excluded persons with diagnoses of osteoporosis or fragility fractures prior to DXA or end of enrollment. Covariates were obtained from the most recent data in the year prior to DXA or, if no DXA, the year prior to enrollment end. Absolute 10-yr hip fracture risk was calculated via FRAXTM, fall risk was estimated using a modified Falls Risk Assessment Tool (FRAT), and comorbidity was assessed via Quan score. Relative risks were estimated from Poisson regression. **Results:** Screening DXA prevalence was 43% (10,061/23,479). Unscreened women were older (77.0 $\pm$ 7.3 yrs vs 72.3 $\pm$ 5.7; mean $\pm$ SD) and had higher comorbidity (4.2 $\pm$ 3.4 vs 3.8 $\pm$ 2.9), FRAXTM hip fracture risk (6.9 $\pm$ 6.9 vs 4.2 $\pm$ 4.5), and fall risk (1.0 $\pm$ 0.9 vs 0.6 $\pm$ 0.81) [all $p < 0.05$]. Multivariate modeling demonstrated lower DXA likelihood for: each added year of age (0.945; 0.042-0.948); each FRAT point (0.765; 0.747-0.783); Blacks (RR: 0.821; 95%CI: 0.756-0.892) and Hispanics (0.892; 0.837-0.950) vs Caucasians; and BMI $\leq$ 18.5 kg/m² vs all other categories (0.763; 0.756-0.892). FRAXTM was not associated with DXA in adjusted analyses and each Quan point increased DXA likelihood (1.030; 1.025-1.036). **Discussion:** The only variable associated with screening DXA that is also known to predict hip fracture risk was Caucasian race. Older age, being underweight and increased fall risk lowered the chance of DXA, but all these factors are known to increase hip fracture risk. Multivariate modeling found greater comorbidity associated with screening, which may represent increased health care utilization and more preventive care opportunities. Further studies should determine whether patient preferences, provider practice patterns, or both underlie the neglect of screening DXA in women at high fracture risk.

Keywords: Screening; Osteoporosis; Prevention and Wellness

doi:10.3121/cmr.2012.1100.cb8-02

CB8-03:

Models of Integrated Care for Adolescent Alcohol and Drug Use in Pediatrics: Predictors and Implications for Practice and Policy

Stacy Sterling¹; Andrea Kline-Simon¹; Charles Wibbelsman¹

¹Kaiser Permanente Northern California

Background/Aims: We describe findings from a longitudinal study of adolescent substance use (SU) treatment, a web survey of pediatric primary care providers (PCPs), and a pilot study of a Screening, Brief Intervention and Referral to Treatment (SBIRT) model of primary care-based adolescent behavioral healthcare. **Methods:** The treatment study (N=419) examined pathways to treatment, co-morbidities, and outcomes. The survey (N=437) examined PCP attitudes and knowledge, patient characteristics, and environmental influences, (e.g., mental health parity and medical marijuana laws), and from electronic medical records (EMR), we examined patient demographics, comorbidity, and services utilization. We examined how PCP,

panel, and organizational characteristics influence screening practices. The pilot examined whether SBIRT versus usual care increased problem identification and specialty treatment rates, and the feasibility of SBIRT in Pediatrics. **Results:** Intakes had high levels of medical and psychiatric comorbidities, and frequent primary care visits, but few had SU problems identified or were referred for treatment by PCPs. Integrated treatment produced better short- and long-term outcomes. The survey found that PCPs were less concerned about alcohol and marijuana than other drug use, rated alcohol use as more difficult to discuss (19% v s 15%) or diagnose (56% vs. 70%) than depression, and were more comfortable discussing sexual practices than alcohol (32% vs. 22%). They were more likely to screen boys than girls, with male PCPs even more likely: 23% vs. 6% ($p < 0.0001$). Self-reported screening rates were far higher than actual (EMR-documented) rates for all substances. Experience, specialty, and recent AOD training (all $p < 0.05$) predicted self-reported rates; only patient age predicted actual rates. Organizational approaches (e.g., EMR tools and workflow guidelines) may matter more than PCP or patient characteristics in determining adolescent alcohol and drug use screening practices. The SBIRT model of care tested in the pilot proved highly feasible. PCPs said that it improved care; more (77) teens were identified and referred for further assessment, and specialty treatment initiation increased from 8.73% to 12% ($p < 0.0001$). **Discussion:** Organizational factors, lack of training, and discomfort with screening may impact adolescent screening and intervention. We discuss the development of integrated models of care for adolescent behavioral healthcare.

Keywords: Integration; SBIRT; Prevention and Wellness

doi:10.3121/cmr.2012.1100.cb8-03

CB8-04:

"I Got Mine, Have You Gotten Yours?" Email to Promote Colorectal Cancer Screening Among Members of a Social Network

Sarah Cutrona¹; Terry Field²; Andrew Williams³; Douglas Roblin⁴; Sarah Greene⁵; Bridget Gaglio⁶; Joann Wagner²; Paul Han⁷; Mary Costanza¹; Brandi Robinson⁴; Vinutha Vijayadeva³

¹University of Massachusetts Medical School; ²Fallon Community Health Plan; ³Kaiser Permanente Hawaii; ⁴Kaiser Permanente Southeast; ⁵Group Health Cooperative; ⁶Kaiser Permanente Mid-Atlantic, MAPMG; ⁷Maine Medical Center

Background/Aims: Members of a social network can influence screening behavior, but it is unclear whether people are willing to share colon cancer screening experiences by email as a means of promoting screening in friends and family. **Methods:** We conducted interviews with 140 people ages 40-70 years in 3 states (Massachusetts, Hawaii, Georgia) to assess their willingness to encourage colorectal cancer screening among friends and family by sharing their own screening experiences. Preferred mode of message transmission (spoken, postcard or email), message characteristics, and projected number of email or postcard recipients per respondent were ascertained. The participants were asked to estimate the impact of their message on their friends or family with regard to willingness to discuss colon cancer screening with a healthcare provider. They were also asked to assess how they would feel receiving such a message. **Results:** 84% of all respondents indicated willingness to pass along a self-edited message in which senders stated that they had completed a colonoscopy and encouraged colon-cancer screening in friends and family over age 50. Among those who report having had colonoscopies, 83% indicated willingness to transmit a message, with 60% indicating they would send emails. Respondents who self-identified as being sources of health information for friends or family were more willing to transmit messages (89.7% willing vs 65.5%; $p < 0.05$). 116 respondents reported willingness to send out an estimated total of 577 postcards or 993 emails. 69% of all respondents thought that receiving such a message themselves would make them more likely to discuss colon cancer screening with a healthcare provider and 64% thought it would make friends and family more likely to have such a discussion. **Discussion:** The majority of interviewees who had completed colonoscopies were willing to share their screening experiences with friends and family through a self-edited email; close to two-thirds thought such an email would influence recipients. Further work is necessary to explore the impact of such an intervention.

Keywords: Cancer Screening; Social Network Communication; Prevention and Wellness

doi:10.3121/cmr.2012.1100.cb8-04