pragmatic design to evaluate health systems-based approaches to CRC screening among age-eligible patients who receive care at FQHCs. **Methods:** STOP CRC has two phases. In Phase I (Year 01), we will pilot-test an evidence-based approach to improve participation in CRC screening in two FQHCs. This phase will include an assessment of the intervention's feasibility, effectiveness, and cost. In Phase II (Years 02–05), we will conduct a comparative effectiveness pragmatic clinical trial, using a mixed-methods approach to evaluate the adoption, implementation, and maintenance of our CRC screening program. Throughout the project, we will involve a diverse planning advisory group of FQHC clinicians and patients, community representatives, state policy makers, and researchers, using principles of Community-Based Participatory Research. **Conclusions:** STOP CRC is a large-scale approach to raising rates of colorectal cancers screening in FQHCs. STOP CRC is expected to expand meaningful use of electronic health information; synergistically meet national goals to improve cancer screening rates, address health disparities, reduce health care costs; and act as a catalyst for future systems-based approaches to improve care delivery in FQHCs and other health systems.

Keywords: Methods; Pragmatic Trial; Colon Cancer

doi:10.3121/cmr.2013.1176.ps1-55

PS1-56:

Beyond Barriers: Systemic Constraints Limiting Sexual Health Care for Breast Cancer Survivors

Meghan Halley¹; Suepattra May¹; Katharine Rendle¹; Dominick Frosch¹; Allison Kurian²

¹Palo Alto Medical Foundation for Healthcare, Research and Education;

²Stanford University

Background/Aims: Sexual health problems represent one of the most frequently experienced and longest-lasting effects of breast cancer treatment, but research suggests that providers rarely discuss sexual health with their patients. Existing research examining barriers to addressing the sexual health concerns of cancer patients has focused on discrete characteristics of the provider-patient interaction without considering the broader context in which these interactions occur. Drawing on focus group discussions with breast cancer survivors, we explore how foundational cultural and structural characteristics of the healthcare system may be preventing breast cancer survivors from addressing their sexual health concerns. **Methods:** Five focus groups were conducted with breast cancer survivors receiving support services at a breast cancer advocacy and resource organization in Northern California. Each group focused on a different aspect of treatment including: 1) diagnosis; 2) surgery and reconstruction; 3) chemotherapy; 4) radiation; and 5) survivorship. An interview guide for each topic area was used to elicit participants' thoughts, opinions and experiences of breast cancer treatment. Analysis utilized inductive techniques incorporating elements of Grounded Theory to identify salient themes that emerged in the discussions. **Results:** An average of eight women participated in each focus group, and women were allowed to participate in more than one group, for a total of 21 participants. Participants' discussions illustrated three core ways in which cultural and structural characteristics of the healthcare system prevented them from addressing their sexual health concerns, including: 1) the structure of cancer care led to participants being disconnected from the healthcare system at the time when sexual side effects most commonly emerged; 2) when their sexual side effects did emerge, the highly specialized structure of the biomedical system made it difficult for patients to identify the appropriate provider to address their complex sexual health concerns; and 3) when patients did discuss sexual health with their providers, their providers approached sexuality as primarily physical, while participants were experiencing biopsychosocial sexual concerns. **Conclusions:** These results suggest that addressing breast cancer survivors' sexual health needs will require a comprehensive approach that includes educating physicians, coordinating existing resources, and developing new sources of information and support during the survivorship period.

Keywords: Breast Cancer; Sexuality; Communication

doi:10.3121/cmr.2013.1176.ps1-56

PS1-60:

Advanced Pancreatic Cancer: Patterns of Care and Recommendations for Education and Support

Jessica Engel¹; Darlene Plank¹; Adedayo Onitilo¹

¹Marshfield Clinic / Security Health Plan of Wisconsin

Background/Aims: Advanced pancreatic cancer has a poor overall prognosis. Fear, anxiety, sadness, and uncertainty are common responses felt by many who are diagnosed with this unfortunate disease, as well as feeling a variety of discomforts related to the cancer itself. Faced with limited time, many hope for longevity while maintaining as good a quality of life as possible. After diagnosis, a multitude of considerations are thought to be essential and beneficial for the patient, caregivers, and medical care providers, including addressing advanced illness planning related to coping, decision-making, cancer treatment, symptom management, and end-of-life care. While the recognition of those considerations is thought to contribute to improved quality of care, often these topics or related interventions are not fully addressed or implemented, potentially leading to a less than ideal experience for the patient with advanced pancreatic cancer. The goal of this study is to evaluate care patterns for advanced pancreatic cancer, address completeness of care based on current standards and guidelines, and provide recommendations for improved overall care and support. **Methods:** Data from the Marshfield Clinic electronic medical record and cancer registry will be electronically and manually abstracted. All patients diagnosed with stage IV pancreatic cancer from 2010-2012 will be included to reflect current care patterns. Data will include patient and cancer characteristics; cancer treatment; presence and management of significant symptoms; number and reason for hospitalization(s); whether or not prognosis and treatment options were discussed; and whether palliative/hospice care and other supportive care referrals were offered or occurred. Comprehensive review of the literature will be undertaken to compile present standards and guidelines regarding advanced cancer care. **Results:** Planned analyses include summary of patient and cancer characteristics, including survival; and determination of variability and commonalities in patient care coordination and utilization of supportive services. **Conclusions:** Understanding the met and unmet needs of patients with advanced pancreatic cancer will lead to improved quality of care and quality of life. Results of this study will be used to write a guideline for improved care and support.

Keywords: Advanced Cancer Planning; Pancreatic Cancer; End of Life Care

doi:10.3121/cmr.2013.1176.ps1-60

PS1-61:

Race and Receipt of Radiation Therapy with Curative Intent after Radical Prostatectomy for Prostate Cancer

Kathryn Richert-Boe¹; Reina Haque²; Stephen Van Den Eeden³; Sheila Weinmann¹; Bhaskar Kallakury⁴

¹Kaiser Permanente Northwest; ²Kaiser Permanente Southern California;

³Kaiser Permanente Northern California; ⁴Georgetown University Medical Center

Background/Aims: We determined if differences existed in prostate cancer treatment received by Caucasian and African-American men who had local or biochemical recurrence following radical prostatectomy (RP) for prostate cancer at an HMO where access to medical care is theoretically equal for all members. We used data collected for a population-based case-control study designed to evaluate the role of specific molecular markers in prostate cancer survival. The subjects were diagnosed with prostate cancer from 1971 through 2001. **Methods:** Study subjects were members of one of three health plans: Kaiser Permanente Northwest (KPNW), Kaiser Permanente Southern California (KPSC), or Kaiser Permanente Northern California (KPMC). The study population included 90 African-American and 233 Caucasian men with prostate cancer who had either local or biochemical recurrence following RP. We compared the likelihood, according to race, of treatment with salvage radiation therapy with curative intent (SRT) among these subjects. We reviewed medical records of all subjects to determine all treatment received in addition to RP and to assess potential confounding factors and effect modifiers. **Results:** After adjusting for case/control status, institution, age at diagnosis, year of diagnosis, Gleason grade (all matching criteria in the