Results: MIST's precision was 0.93 and recall was 0.77. MIST left 13 leaks and incorrectly identified two instances of blood pressure numbers as dates. We can reduce 3 out of 13 leaks using structured data obtained for that visit note. The remaining leaks are: 3 locations, 1 age which belonged to the patient's child, 1 visit date pre-EHR system, 3 pieces of historical visit data, and 2 mentions of 'sheriff' because the chart abstractor determined it to be an "other" PII type. **Conclusions:** Our results show that combining structured data with MIST can potentially improve de-identification of Group Health clinical text. Future work is to build an automated system that include historical visit data, use gender, race, and language to create more realistic surrogates and to evaluate the loss of important clinical information in de-identified documents.

Keywords: Natural language processing; De-identification
doi:10.3121/cmr.2014.1250.c3-4

PS1-45:

Get Smart: Finding Structured Data Using SmartTools in Clarity

Sharon Fuller¹

¹Group Health Research Institute

Background/Aims: Chart notes and other text blocks in Epic's electronic medical record system generally appear to be free text, requiring natural language processing (NLP) for analysis. In fact, though, many notes contain boilerplate text that can be traced back to structured data created using various Epic "SmartTools." We investigated Clarity, Epic's reporting database, to learn how to locate these structured data. **Methods:** Through trial and error we developed a data map to guide searches for structured data in note text. Starting with a known block of text, we found the underlying boilerplate, traced the sources of the individual data elements, and finally located the values recorded for those elements in the encounter. **Results:** Smart phrases, lists, links and forms are a few of the commonly used Smart elements in Epic. Each of these elements may be composed of other elements, increasing the difficulty of charting a straight path. So a SmartPhrase may include the option to choose from a SmartList, and those choices may, in turn, reference SmartLinks. However, if one can locate, for example, a definitive statement of pathology results within the structured Smart data of a results letter to the patient, it may be possible to avoid much of the expense and uncertainty involved in using NLP to parse the same information. Similarly, if some note text is known to be a direct copy of structured data from another part of the chart, one can safely exclude that text from further examination. Even if one does not analyze the structured data directly, discovering which text is "Smart" and the expected values for the included data elements can greatly facilitate the development of NLP algorithms. It is important to work with clinicians at this first level of discovery, to learn the mnemonics or "dot-phrases" they use to invoke SmartTools and the text that results. **Conclusions:** Epic's SmartTools hold great promise for teasing structured data out of apparently unstructured text. However, unravelling the complex interrelations among SmartTools requires a good deal of motivation and ingenuity. Further investigation is needed to determine the most appropriate and effective uses of these data.

Keywords: Clarity; Structured data
doi:10.3121/cmr.2014.1250.ps1-45

PS1-13:

Probabilistic Linkage (Also Known as "Fuzzy Matching"): the Theoretical Foundations of Modern Record Linkage

Christopher Mack¹

¹Group Health Research Institute

Background/Aims: Data joins have historically been done using deterministic linkage techniques which require exact data matches to succeed. For data extracted from disparate sources, requiring an exact match may be too limiting. Probability-based linkage techniques have gone from an arcane art involving expensive third-party software to having multiple free and commercial options available for use. We will walk through the theoretical underpinnings of the probability-based linkage theory used by many record linkage applications today. **Methods:** Describe the history of record linkage, the cause of linkage errors, the need for probability-based

techniques for match assessment and the mathematical framework of probabilistic linkage. Discuss matching best practices and limitations in matching ability and match assessment. **Results:** The audience will better understand how to choose linking variables, ways to limit the search space for comparisons, and the importance of pre-processing linkage variables, including comparison by permutation of variables, phonetic codes and edit-based measures. **Conclusions:** The audience will be able to better evaluate, and to use more successfully, the available freeware and commercial linkage software given a better understanding of their inner workings, limitations and potential pitfalls. An advanced audience will be in a better position to create their own probabilistic matching algorithms.

Keywords: Linkage; Matching

doi:10.3121/cmr.2014.1250.ps1-13

PS1-14:

Pregnancy Data in Clarity - Stay Tuned for Our Next Episode

Sharon Fuller¹

¹Group Health Research Institute

Background/Aims: For a condition that is so easily defined, pregnancy can be remarkably difficult to locate with any precision in existing structured data sources, especially while a woman is still pregnant. Researchers may require pregnancy information: (a) to avoid recruiting pregnant women to clinical studies; (b) to recruit women in a certain trimester or meeting certain pregnancy-related criteria; (c) to exclude measures taken during pregnancy; (d) to evaluate prenatal care across populations; (e) to provide conditional variables for other analyses, including neonatal outcomes. In 2004, Group Health (GH) adopted Epic's electronic medical record system. In this poster, we investigate Clarity, Epic's reporting database, to determine its usefulness as a source of structured pregnancy data. **Methods:** We examined all OB episodes of care at GH clinics that included any part of 2012. We also examined records of all completed visits in 2012 with an encounter type of OB Visit. **Results:** Almost all OB visits recorded in Epic at GH are part of an OB episode and pregnancy dating information is almost universally recorded. OB episodes assemble a large amount of useful data, with more variables anticipated following the recent implementation of Epic 2012 at GH. It remains to be seen how many of these will be populated, and if any will be backfilled from other sources. Although prenatal episode data are quite complete, parity, delivery and outcomes information at GH require more complex querying and result in less definitive data. Nor are episode data available for patients who obtain prenatal care outside of the GH network of clinics. Episodes may also be useful for examining the overall care picture for other health conditions. **Conclusions:** OB episodes in Clarity can be a rich source of structured data for current and past pregnancies, although implementations of Epic and Clarity at different sites may vary in the specific data elements captured.

Keywords: Pregnancy; Clarity

doi:10.3121/cmr.2014.1250.ps1-14

PS1-15:

Pre-filling Breast MRI Abstraction Forms Using Natural Language Processing

Hongyuan Gao¹; Karen Wernli¹

¹Group Health Research Institute

Background/Aims: Information in breast MRI reports is valuable for breast cancer research, but these data are only available in free-text reports and require resource-intensive manual abstraction. We developed and tested a Natural Language Processing (NLP) algorithm to extract information and pre-fill abstraction form from free-text breast MRI reports. **Methods:** We identified 465 reports for women receiving breast MRI at Group Health between 2010-2012. We developed an NLP algorithm in SAS v9.2. The algorithm extracts information of reading radiologist, laterality, parenchymal enhancement, whether computer-aided technique is used, comparison exams, clinical indications and assessment from breast MRI reports. The NLP results are compared with manual abstraction from an experienced abstractor. **Results:** The algorithm correctly extracts reading radiologist, laterality and whether computer-aided technique for all 465 breast MRI reports, except 1

report with inconsistent information on laterality itself. It correctly extracts 83% of 465 reports for assessment for right breast and 92% for assessment for left breast. Unstable gold standard impedes performance of the NLP algorithm for extracting parenchymal enhancement and clinical indications. There is no gold standard to show NLP performance for comparison exams yet. **Conclusions:** This NLP algorithm holds promise for rapid, accurate extraction of information from free-text breast MRI reports. Manual review will be faster and more accurate due to the pre-filling of the abstraction form. **Keywords:** Natural language processing; Breast MRI report

doi:10.3121/cmr.2014.1250.ps1-15

PS1-16:

Effect of Insurance Plan Choice on Utilization in Kaiser Permanente Colorado

Nikki Carroll¹; Laurie Crounse¹; Jeffrey Holzman¹; Carsie Nyirenda¹

¹Kaiser Permanente Colorado

Background/Aims: There has been a shift in recent years from traditional HMO-type insurance products to high deductible or co-insurance plans. Kaiser Permanente Colorado (KPCO) has also added new service areas to the region that include Southern and Northern Colorado. In addition, the impending Affordable Care Act will be implemented in 2014 which may bring new and different members to KPCO. There is a need to describe socio-demographic and clinical characteristics by plan type in order to understand how members within these different plan types utilize KPCO resources and what data are available and identifiable within our source data systems. These answers will enable us to establish appropriate inclusion and exclusion criteria for future research studies. **Methods:** All members with at least 1 day of membership from January, 2000 through July 2013 were identified. Membership within product lines and all associated utilization was based on calendar year. Patient characteristics, ambulatory visits, inpatient admissions, emergency department admissions, lab tests, tumors, pharmacy utilization and variations in therapy for certain diagnoses were obtained from the Virtual Data Warehouse (VDW). Socio-demographic characteristics, including gender, race, and age, were described by percentages. Clinical characteristics and variation in utilization were calculated on a per-person per-month measure and were compared for each region/insurance product/year combination. **Results:** Demographics and utilization varied across all product lines and years. An increase in the number of different products offered was observed over time. Members in the HMO plan type were more likely to be younger, less educated and use more resources than members in other products. Some data were not easily identifiable in certain plan types. **Conclusions:** These results indicate that demographic and clinical patterns do vary between plan types. These patterns need to be taken into account when creating cohorts in future research projects.

Keywords: Utilization; Insurance plans

doi:10.3121/cmr.2014.1250.ps1-16

PS1-17:

Organization-wide Collaboration to Convert Binary Clinical Notes to Research-ready Text

Scott Halgrim¹; Shane Reeh²; Roger Kelem²; Roger Wong²; Aruna Kamineni¹; Virginia Immanuel¹; Christopher Pattillo²; Roy Pardee¹; David Carrell¹; Carolyn Rutter¹

¹Group Health Research Institute; ²Group Health

Background/Aims: Clinical text is an integral part of research at Group Health Research Institute (GHRI), supporting ongoing projects and new research proposals. Our clinical notes are stored in a single, full-text-indexed table that is populated nightly with all new notes in Group Health's Electronic Medical Record (EMR). These notes include those generated internally and those received from partner organizations via electronic interfaces. Recently one of our partners changed its interface from sending formatted text to sending binary PDF files. While this had little effect on clinical users, it made a significant amount of text data inaccessible for computerized processing. This text was crucial to ongoing research, so we needed to find a way to preserve access. **Methods:** A diversely-skilled team of technologists collaborated to solve the problem with the following

process. When the nightly HL7 file arrives at Group Health, it is copied to a research server and the message ID is stored with the appropriate encounter in the EMR. A scheduled Python script then extracts the message ID and binary PDF from each HL7 message. It converts the base64-encoded information into a PDF file and names it with the message ID for linking with the encounter. Finally, the script converts the PDF file into a text file using an open source library. The PDF and text files are archived, as is the nightly HL7 message. **Results:** From June to October 2013, this process has converted and archived over 30,000 notes from our partner that otherwise would not have been available for research. The files are also available to Group Health's claims auditing department, enabling Group Health to see a monetary return on the effort. **Conclusions:** Thanks to the open source community and strong connections between informatics management at Group Health Cooperative and GHRI, we were able to quickly salvage critical data for research when an externally-driven change to that information occurred. Before the end of 2013 we expect to integrate these converted text notes with our clinical text storage.

Keywords: Natural language processing; Data interfaces

doi:10.3121/cmr.2014.1250.ps1-17

PS1-18:

Do Patients Use Online Messaging to Substitute for Office Visits and Phone Calls? Results from a Natural Experiment

Laura Panattoni¹; Sean McClellan¹; Ming Tai-Seale¹

¹Palo Alto Medical Foundation Research Institute

Background/Aims: We evaluated the effect of online messaging between patients and providers on other healthcare utilization. Messaging may increase the administrative burden on providers but may also substitute for utilization such as calls or visits. To identify the effect of messaging, we exploited a natural experiment, which occurred in 3/2011, when a large multispecialty practice in California lowered the price of messaging from \$60/year to free for patients and initiated incentives for doctors (\$3/thread), resulting in increased rates of patients messaging. **Methods:** We conducted a longitudinal study of 65,332 active primary care patients from 3/2009-3/2012. All patients in the study had activated their online portal before 3/2011 and none had paid to use messaging previously; 38,438 of these patients initiated a message following 3/2011. Messaging was measured by the number of threads initiated by patients annually. Outcome measures included annual rates of office visits and phone calls to primary and specialty care. The effects of messaging were identified through difference-in-differences analyses using generalized least squares, comparing the utilization of those initiating any messages with those not messaging, before and after messaging became free. Controls included patient fixed effects along with patient age, insurance type, Charlson score, continuity of care, and relevant characteristics of their primary care provider. **Results:** Comparing the rates of service use before and after 3/2011, between messaging and non-messaging users, we found no difference in the annual number of primary care phone calls (0.002 per patient-year, $P = 0.394$), and a small increase in primary care office visits (0.05 per patient-year, $P < 0.001$) for each thread initiated. For specialty care, we found an increase in 0.16 phone calls ($P < 0.001$) and 0.21 ($P < 0.001$) office visits per patient-year for each thread initiated. **Conclusions:** Among patients who were already using the online portal, use of secure messaging was associated with little difference in primary care utilization, but with a moderate increase in specialty utilization. Although current analyses controlled for patient and provider characteristics, patient self-selection may still threaten validity; we will address this through propensity score stratification in future analyses.

Keywords: Online messaging; Healthcare utilization

doi:10.3121/cmr.2014.1250.ps1-18

Health Services Research/Health Policy/Costs

B2-1:

Estimation of Standardized Hospital Costs from Claims Data that Reflect Resource Requirements for Care

John Schousboe¹; Misti Paudel²; Brent Taylor²; Lih-Wen Mah³; Beth Virnig²; Kristine Ensrud²; Bryan Dowd²