Algorithm for Real-Time Identification of Blood Pressure Data Entry Errors Using Longitudinal Data From the Electronic Health Records

G. Craig Wood, MS, *Geisinger Center for Health Research*; Jason Roy, PhD, *Geisinger Center for Health Research*; Nirav R. Shah, MD, MPH, *Geisinger Center for Health Research*; James B. Jones, MBA, *Geisinger Center for Health Research*; H. Lester Kirchner, PhD, *Geisinger Center for Health Research*; Walter F. Stewart, PhD, MPH, *Geisinger Center for Health Research*

Background/Aims: Data from electronic health records (EHR) are prone to error due to collection methods or human entry. However, historical measures can be used to validate subsequent measures. Real-time algorithms can be created that utilize longitudinal data to identify and require confirmation of questionable data during the patient encounter. **Methods:** We developed algorithms for identifying potential errors in the Geisinger Health System EHR. These were developed with two key principals in mind: (1) the algorithms should be relatively simple to program and implement, and (2) the identification of erroneous values should depend, to varying degrees, on both a patient's historical data and on data from the overall population. The more encounters the patient has had in the past, the more heavily the outlier cutoff values should depend on that patient's historical data. The algorithms were tested and refined on a sample of over 2 million systolic blood pressures from over 212,000 individuals. **Results:** The selected algorithm utilized prior measurements in the EHR and allowed for the possibility that the stability of the measurements might vary across subjects (e.g., the standard deviation might vary with the mean). The properties of the algorithm are described and are used to identify cutoffs for defining 5%, 1%, and 0.1% of the measurements as questionable data. Future studies are planned to test the usefulness of the algorithm in real time. In addition, these studies can be used to identify a cutoff threshold that results in an appropriate balance between flagging too many accurate values, and not flagging enough inaccurate values. **Conclusions:** An algorithm that uses historical blood pressure data can be used to identify potential errors during the patient encounter. The relatively simple algorithm does not require complex programming/software and likely can be tailored to optimize use by clinicians. The implementation of such an approach has the potential to improve the quality of EHR data.

Applications of Mediation Analysis in Behavioral Medicine Research

Douglas W. Roblin, PhD, *The Center for Health Research/Southeast, Kaiser Permanente Georgia*; Jeroan J. Allison, MD, MSc, *School of Medicine, University of Alabama at Birmingham*; Edmund R. Becker, PhD, *Rollins School of Public Health, Emory University*; Peter J. Joski, MSPH, *The Center for Health Research/Southeast, Kaiser Permanente Georgia*

Background: In a typical regression model, the total effect of an independent variable (e.g., social support or strain) on a dependent variable (e.g., health status) is estimated. In mediation analysis, a regression model accounts for the possibility that the total effect consists of a direct effect and an effect mediated through a third variable (e.g., patient activation). We examined the potential mediating effects of patient activation between various psychosocial circumstances and health status, dietary intake, exercise, and obesity among a sample of adults in a group-model managed care organization (MCO). **Methods:** A mixed-mode survey (written and Internet options) was administered from October through December 2005 to 5309 MCO enrollees, 25-59 years of age, who were randomly sampled from 3 condition cohorts: (1) adults with type 2 diabetes, (2) adults with elevated lipids but no acute CVD history, and (3) adults with no major morbidities (low risk). The survey included previously validated measures of psychosocial circumstances that measured support or strain between patients and physicians (care delivery climate), among friends and family members (social climate), and among coworkers (work climate); patient activation (PAM-13); healthy behaviors (e.g., exercise, dietary intake); and physical and emotional health (SF-12). Using the approach of Baron and Kenney, linear and logistic regression models were estimated to compute the mediating effect of patient activation. The mediating effect was tested for significance using the Sobel test. **Results:** Twenty-two hundred twenty-four completed the survey (42% response rate). Forty-five percent of respondents were black; 60% were female. Adjusted for condition cohort and other patient covariates,

psychosocial circumstances tended to have significant ($P<0.05$) total and direct effects on health status (better with more supportive climates), diet and exercise (better with more supportive climates) and BMI (lower with more supportive climates). Patient activation was a significant mediator (Sobel $P<0.05$) of the observed relationships between psychosocial circumstances and each of the three outcomes (health status, health behavior, and BMI) for all cohorts. **Conclusions:** Patient activation has an important mediating role between psychosocial circumstances and health and healthy behaviors, particularly with respect to social climate.

Evidence-Based Medicine Integrator (EBMI): A New Simulation Architecture for Chronic Disease

Jonathan Brown, MPP, PhD, *Kaiser Permanente Center for Health Research*

Background: Computer simulation models are increasingly used in complex chronic disease to prioritize treatments, study the effects of treatments, and guide policy and management. The accuracy of these models is doubtful, particularly in real-world use. **Methods:** We designed EBMI to 'think' as an evidence-based clinician thinks, by combining the best user-specified trial evidence about treatment effectiveness with the best possible individualized estimates of patient risk. Local validity is much more likely because EBMI's risk functions can be estimated from local data. The use of local data also allows the model to be used for real-time clinical prioritization in local settings. **Results:** EBMI reproduces Kaiser Permanente Northwest data very closely and, in its only validation to date against a clinical trial, closely replicated the Heart Protection Study. More validations and studies using the model are underway. **Conclusions:** It is possible to design a simulator that is both more evidence-based and more locally valid than existing models. EBMI and its source code are now available to other researchers, clinicians and managers.

Diabetes

Factors Associated With Continued Inertia Among Patients With Diabetes

Jennifer E. Lafata, PhD, *Center for Health Services Research, Henry Ford Hospital*; Elizabeth A. Dobie, MPH, *Henry Ford Hospital*; George Divine, PhD, *Henry Ford Hospital*; Bruce D. McCarthy, MD, *Allina Medical Clinic, Minneapolis, MN*; Marianne Ulcickas Yood, PhD, *Yale University, New Haven, CT*

Objectives: To estimate the prevalence of and factors associated with inertia among patients with diabetes. **Methods:** Using automated clinical and pharmacy data, we identified an insured cohort of patients with diabetes at the time they initiated oral monotherapy during the period, 2000-2005 ($n=5082$). Actuarial methods were used to estimate time to inertia (defined as 2 HbA1c values over 8% at least 90 days apart, not yet dispensed insulin with no medication change in the preceding 90 days). Among the subset facing inertia ($n=1391$), actuarial methods and Cox regression were used to estimate time to and factors associated with appropriate care defined as either a medication intensification (dose change, class change, or class addition) or an HbA1c $\leq 7\%$. **Results:** At inception, mean age of the cohort was 60.1 years, mean HbA1c was 8.6% and 97% were dispensed either Sulfonylurea or Metformin. Forty-eight percent were female and 37% African American. Eight percent of patients faced inertia within 1 year, 18% within 2 years, 26% within 3 years and 33% within 4 years. Before the time of inertia, 35% had been dispensed combination therapy. At the time of inertia, 20% had no medication on hand (i.e., were non-persistent in their medication use) and mean HbA1c was 9.7% (range 8.1%-18.9%). Mean time to appropriate care was 7.3 months, with 27% having neither an intensification nor HbA1c $\leq 7\%$ within 1 year. Those with no medication on hand (hazard ratio (HR)=0.67; 95% CI, 0.56-0.82) and those who had progressed to combination therapy (HR=0.86; 95% CI, 0.76-0.98) faced longer delays. Increased delays were also associated with increasing drug co-payments (HR=0.95; 95% CI, 0.92-0.98). HbA1c levels greater than 11% (HR=1.48; 95% CI, 1.11-1.97) were associated with shorter delays, as were dispensings for an anti-hypertensive (HR=1.17; 95% CI, 1.01-1.37), lipid lowering (HR=1.19; 95% CI, 1.03-1.37), and anti-depressant (HR=1.46; 95% CI, 1.17-1.82)

medication. Increasing visit frequency to primary care (HR=1.85; 95% CI, 1.58-2.18) and endocrinology (HR=2.08; 95% CI, 1.45-2.97) were associated with decreasing delays, as were increasing income levels (HR=1.04; 95% CI, 1.00-1.07). **Conclusions:** Patients with diabetes facing inertia are at risk of further delays in appropriate management. Our findings suggest the presence of patient, physician and system barriers to appropriate care. Increased contact with the health care system may mitigate risk.

Abstract C-C1-03

Moving the Big N

Ellen Vanderboom, MSN, RN, *Marshfield Clinic*; Theodore Praxel, MD, MMM, FACP, *Marshfield Clinic*; Marilyn Follen, RN, MSN, *Marshfield Clinic*; Daniel Erickson, MD, *Marshfield Clinic*; Thomas Gabert, MD, MPH, *Marshfield Clinic*; Kori Krueger, MD, *Marshfield Clinic*; Eric Penniman, DO, *Marshfield Clinic*; Mary Beth Dickinson, RN, BSN, *Marshfield Clinic*; Trina Ford, MSN, NP, RD, *Marshfield Clinic*

Background: Nationally the trend for hospitalization of diabetics has been slow to show improvement. In 2003, the rate of hospital discharge for diabetes as any-listed diagnosis (360.7 per 1000 people with diabetes) was only slightly lower than the rate in 1980 (398.7 per 1000 people with diabetes). Decreasing hospitalizations and improving health outcomes for 17,678 patients has been a system wide initiative in Marshfield Clinic since 2005. **Methods:** Numerous interventions have occurred since 2005 to improve health outcomes and decrease costs for the diabetes population. Evidenced-based guidelines were deployed and CME activities were created for providers. Clinical information support tools and applications deployed included wireless tablets for primary care providers and staff, an electronic point of care reminder tool, medication management and electronic planned care applications were also deployed. The diabetic foot exam process was standardized and medical assistants were trained to perform foot exams. Standing orders for staff to give influenza and pneumococcal vaccinations and schedule necessary lab tests or office visits were developed. Feedback on diabetes quality metrics at the organization, division, department and individual primary care provider level was provided every quarter on a rolling 12-month basis. **Results:** The all-cause hospitalization rate for persons with a diagnosis of diabetes has decreased from 360 per 1000 in 2005 to 317 per 1000 in 2007. This decrease in hospitalization lead to improved economic outcomes and quality of life by avoided hospitalizations. Conservative estimates of the cost savings range from \$5,202,000 to \$14,137,200. **Conclusions:** Significant improvement in this population has been demonstrated system wide by a decrease in all-cause hospitalization rates and composite diabetes process and outcome measures. Use of an electronic health record is essential but not sufficient to affect change. Ongoing measurement of practice performance has assisted in the change. Physician champions have been crucial for spread. Improvements are significant, however, opportunities remain.

Abstract C-C1-04

The Effectiveness of Diabetes Care Management in a Large, Integrated Delivery System: a TRIAD Study

Julie A. Schmittiel, PhD, *Kaiser Permanente Northern California Division of Research*; Connie Uratsu, BA, *Kaiser Permanente Northern California Division of Research*; Bruce Fireman, MA, *Kaiser Permanente Northern California Division of Research*; Joe V. Selby, MD MPH, *Kaiser Permanente Northern California Division of Research*

Background: Numerous studies have examined the effect of care management (CM) programs on diabetes outcomes such as A1c control. Most evaluations have been small randomized controlled trials; few studies examine how CM translates to real-world settings in terms of success at reaching target populations, or effectiveness for improving outcomes in large populations. The aims of this project are to assess the appropriateness of entry into a diabetes (DM) nurse-delivered CM program in Kaiser Permanente Northern California (KPNC) compared with program guidelines, to determine the proportion who meet program entry criteria, and to compare the trajectory of intermediate outcomes in patients who receive CM to that of similar patients who don't receive CM. **Methods:** KPNC automated databases were used to identify all DM patients from 2000-2005 and to determine whether patients enrolled in CM. We examined whether CM enrollees and non-enrollees met

the eligibility criteria for the CM program. Using propensity score matching, we matched CM patients to similar patients who were eligible but did not enter CM. We compared trajectories of A1c, LDL-c, and blood pressure levels of CM patients to those of their matched controls, adjusting for baseline risk factor levels. **Results:** Between 2000 and 2005 less than 10% of all KPNC diabetes patients entered CM, representing approximately 25% of all eligible patients. Up to 22% of those in CM each year did not meet program entry criteria. CM patients were similar to the overall DM population in age, gender, blood pressure, and LDL-c levels; they had higher A1c and microalbumin levels. There were small but statistically significant differences in A1c, LDL-c, and blood pressure levels between CM patients and their matched controls 6, 12, and 15 months after program exit. There were significant improvements in CVD risk factor levels in all KPNC DM patients over time, regardless of whether or not they received CM. **Conclusions:** While eligibility guidelines for diabetes nurse care management were not strictly adhered to, care management did have a small but significant impact on intermediate outcomes. However, these improvements were small compared to the risk factor improvements over time in the overall DM population.

Abstract C-C1-05

Who Does Not Complete a First-Fill for a Diabetes-Related Prescription?

Nirav R. Shah, MD, MPH, *Geisinger Center for Health Research*; Annemarie G. Hirsch, MPH, *Geisinger Center for Health Research*; G. Craig Wood, MS, *Geisinger Center for Health Research*; Amanda F. Wingate, BS, *Geisinger Center for Health Research*; Ilene G. Ladd, MS, *Geisinger Center for Health Research*; Walter F. Stewart, PhD, MPH, *Geisinger Center for Health Research*

Background: Research on medication adherence is primarily conducted using insurance or pharmacy claims data, focusing on patients who have already obtained their first prescription for a defined condition. Claims data alone cannot be used to assess first-fill rates. **Methods:** We conducted a retrospective cohort study that examined first-fill rates for patients prescribed medications to treat diabetes by linking prescribing information from electronic health records (EHR) to pharmacy claims data of one insurer. EHR data were obtained from the Geisinger Clinic, a large group practice serving a 31-county area of central and northeastern Pennsylvania. The Clinic includes 41 community practice sites with primary and multi-specialty care, all of which have used an EHR since 2001. Claims data were obtained from Geisinger Health Plan, a payer that accounts for 30% of Geisinger Clinic patient volume. Adherence was defined as a claim generated for the first-fill prescription within 30 days of the index date. **Results:** Of 1132 patients written a new, first-time prescription for any diabetes medication, 962 (85%) generated a corresponding claim within 30 days. Therapeutic class, refills, co-pay, baseline A1c values, and change in A1c values were associated with first-fill rates. Gender, race, age, comorbidity score, number of office visits, and number of medications prescribed had no association ($P < 0.05$). **Conclusions:** Adherence relates directly to higher A1c levels and is associated with therapeutic drug class, higher number of refills, and lower co-pay. As expected, patients with higher adherence exhibited greater change in A1c levels. Patients with relatively lower baseline A1c levels, while at lower risk for diabetes complications, are at higher risk of not adhering to diabetes medications.

Abstract C-C1-06

Identifying Depression Among Diabetes Patients Using Natural Language Processing of Office Notes

Lucy R. Fischer, PhD, *HealthPartners Research Foundation*; William A. Rush, PhD, *HealthPartners Research Foundation*; John C. Kluznik, PhD, *HealthPartners Research Foundation*; Patrick J. O'Connor, MD MPH, *HealthPartners Research Foundation*; Ann M. Hanson, BS, *HealthPartners Research Foundation*

Purpose: To compare two methods for identifying depression among health plan enrollees with diabetes—administrative data base codes versus natural language processing (NLP) of office note verbal narratives. **Background:** In research studies in health care systems, depression cases can be identified through coded data in the health plan administrative data system, including diagnosis (ICD-9) codes and pharmacy codes. However, depression is often