monitoring of this vaccine's safety in early fall 2009. **Methods:** The VSD Project conducts near real-time monitoring of potential adverse events following vaccination for all new vaccines. Each week, each site updates dynamic data files, and the coordinating center analyzes all data cumulatively. Repeated analysis of the same data requires adjustment for multiple testing. To address this issue, VSD uses sequential statistical methods. Evaluation of H1N1 vaccine safety poses special challenges, including confounding by indication, seasonality, and new priority groups including pregnant women. We are using several analytic methods, including: (1) Self-controlled case-series, which compares exposed vs. unexposed time windows within the same individual; (2) Poisson, which compares the outcomes during the current season with expected rates based on historical data for seasonal influenza vaccine; and (3) Difference-in-difference, which compares the relative risk in exposed vs. unexposed time windows for H1N1 vaccine during the current season with the relative risk for seasonal influenza vaccine during cumulative prior seasons. **Results:** Pre-specified outcomes of interest include: Guillain-Barre syndrome, neuropathies, seizures, encephalitis, Bell's palsy, ataxia, anaphylaxis, spontaneous abortion, pre-eclampsia, stroke, myocarditis and wheezing. Historical rates of these outcomes were calculated for time windows after seasonal influenza vaccination in prior seasons (2000/01-2008/09). The sequential analysis methods above will be used, and relative and absolute risks will be estimated for each outcome. For outcomes found associated with vaccination in preliminary analyses, additional evaluation will be conducted using temporal scan analyses, logistic regression, and when appropriate, medical record review.

Keywords: H1N1 safety, Influenza A vaccine monitoring, Vaccine Safety Datalink Project

doi:10.3121/cmr.2010.943.c-b5-02

C-B5-04:

An Algorithm for Classifying Potential Cases of Amyotrophic Lateral Sclerosis from Electronic Health Records Using Decision Tree Analysis

Kathleen Albers, MPH¹; Stuart Hoffman, DO²; Judith Readon, MS²; Mark Hornbrook, PhD³; Alison Naleway, PhD³; Joanna Bulkley, PhD³; Susan Andrade, PhD⁴; Daniel Newman, MD⁵; David Nerenz, PhD⁵; Margaret Gunter, PhD⁶; James Burmester, PhD⁷; Priscilla Velentgas, PhD⁸; Daniel Strickland, PhD⁹; Stephen Van Den Eeden, PhD¹⁰

¹Kaiser Permanente; ²Geisinger Medical Center; ³Kaiser Permanente Northwest; ⁴Meyers Primary Care Institute; ⁵Henry Ford Health System; ⁶LCF Research; ⁷Marshfield Clinic Research Foundation; ⁸Harvard Pilgrim Health Care; ⁹Kaiser Permanente Southern California; ¹⁰Kaiser Permanente Northern California

Background/Aims: The CDC is developing a national amyotrophic lateral sclerosis (ALS) registry that would be based on electronically available medical records and claims. The HMORN contracted with the CDC to help examine the feasibility of doing this in locations (e.g., HMOs) where complete longitudinal records are more likely to be available. Any final system used by the CDC will require an algorithm to classify putative cases. Our aim was to develop an automated algorithm to classify putative cases of ALS and other motor neuron diseases using electronic health records.

Methods: We identified potential cases of ALS by review of inpatient, outpatient, other provider services, mortality and pharmacy data for any evidence of ALS or, more generally, motor neuron disease (MND) from 2001 to 2005 at Kaiser Permanente, Northern California. The records of each individual were reviewed to classify into five categories: probable ALS, not ALS, other MND (not ALS), insufficient information, or unclear. Classification And Regression Tree (CART) analyses were used to create an algorithm based on a comprehensive set of variables that may predict the likelihood of having true ALS (or, conversely, ruling them out). **Results:** Multiple algorithms were developed with key variables being number times an ALS code appeared in the record, physician-type making diagnosis, alternative diagnoses, mortality records, and length of follow-up. In our best model, we were able to correctly classify 99.6% of those originally determined to be probable ALS. However, there remain a substantial number of individuals classified by review or algorithm in the unclear, insufficient, or other MND categories. In addition, with this analytic approach, the "costs"

of misclassification can be modified, and we will present alternative algorithms with a discussion of the benefits and disadvantages of each. Validation analysis results conducted in a second HMO will be presented. **Conclusions:** An algorithm was developed that appropriately classifies the vast majority of probable ALS cases. However, given the lack of detail in some records and the difficulty in diagnosing some clinical cases, there will be some misclassification when based solely on electronic medical records.

Keywords: Amyotrophic Lateral Sclerosis, Electronic health records, ALS Classification

doi:10.3121/cmr.2010.943.c-b5-04

PS1-15:

A Method for Discovering Variant Spellings of Terms of Interest in Clinical Text

David Carrell, PhD¹; David Baldwin, BS¹

¹Group Health

Background and Aims: Mining information from unstructured clinical text requires knowledge of synonyms and variant spellings corresponding to the term of interest. Lexicons such as the Unified Medical Language System identify recognized synonyms but are less helpful in identifying variant spellings or locality-specific language. Natural language processing algorithms employing dictionary look-ups frequently miss terms because of variant spellings, accounting for as much as 25% of algorithm failures. This problem is not easily resolved deductively or through domain expertise. We describe an inductive, empirically-driven method that identifies synonyms/variant spellings of a concept of interest. **Methods:** The conceptual basis for this method is the principle that terms with similar meanings tend to be found in similar textual contexts. Accordingly, sets of words found with high frequency in the context of two synonyms will overlap with high probability. Call the word of interest the target term; call the high-frequency words around it context terms; call words most likely to be synonyms/variants synonym candidates. A four-step process can be used to identify variants. Step 1: Identify the target term's highest frequency context terms. Step 2: Identify terms found in high frequency near each of the context terms. Step 3: Eliminate Step 1 & 2 terms appearing more often outside the context than in it. Step 4: Manually review remaining synonym candidates to identify true synonyms/variants. Using SAS we illustrate the method for target term HER2/NEU in Group Health chart notes for 2008, representing 200,913,281 lines of clinical text. **Results:** Four of the top 20 context terms near HER2/NEU had ratios of in_context_occurrence/out_of_context_occurrence > 1: OVEREXPRESSION, ONCOGENE, OVEREXPRESSED, C-ERBB-2. These context terms occurred 1196, 617, 540, and 170 times, respectively, and produced lists of 205, 119, 143, and 54 of their own context terms, respectively (excluding words with frequency < 2). Manual review yielded a final set of 23 synonyms and spelling variants, including HER-2, HER/2, HER-2-NEU, HER2/NUE (note spelling: "NUE"), FISH, IHC, and C-ERBB-2 among others. **Conclusion:** The method described provides an easily implemented approach to identifying synonyms/variants of terms of interest that can be used to enhance strategies for identifying concepts in clinical text.

Keywords: Identifying variant spellings, clinical text, medical terminology
doi:10.3121/cmr.2010.943.ps1-15

PS1-16:

Disenrollment Rates Among HMORN Sites by Age, Gender, and Medical Condition

Jennifer Ellis, MBA, MSPH¹; Marsha Raebel, PharmD^{1,2}; Addie Clark, BA¹; Susan Andrade, ScD³; Robert Davis, MD, MPH⁴

¹Kaiser Permanente Colorado; ²University of Colorado; ³Fallon/Meyers; ⁴Kaiser Permanente Georgia

Background: It is often desirable to include enrollment and disenrollment (or, conversely, retention) information in grant applications. Overall disenrollment has been evaluated as part of Virtual Data Warehouse quality