colonoscopy “(42%) followed by “I do not think I need the test” (21%). Males were more likely than females to report that they did not think they needed the test ($p = 0.042$) and that they could not take the time off of work ($p = 0.025$), and that they had a bad experience with colonoscopy ($p = 0.027$). **Conclusions:** Survey responses point to the important roles healthcare providers and approaches tailored to the barriers to specific groups can plan in intervention strategies.

Keywords: Colorectal cancer screening, Colonoscopy barriers, Intervention strategies

doi:10.3121/cmr.2010.943.ps3-10

PS3-12:

Obesity and Risk of Proliferative Hyperplastic Lesions

Melissa Stahl, MPH¹; James Evans, MD, MPH¹; Azadeh Stark, PhD¹

¹Geisinger Health System

Background and Aims: Obesity is a risk factor for breast cancer for postmenopausal women. Aromatization of androstenedione to estradiol in the peripheral adipose tissue and low levels of sex-hormone binding globulin as well as increased serum levels of insulin-like growth factors have been proposed as the underlying biological mechanisms. Women diagnosed with hyperplastic lesions with/without atypia have increased risk for a subsequent breast cancer. We conducted a retrospective study with the primary objective of assessing the association between body weight and the diagnosis of hyperplastic lesions. Our specific aims were: 1) to compare the prevalence of biopsies prior to the diagnosis of breast cancer and 2) to estimate the risk of hyperplastic lesions after adjusting for Body Mass Index (BMI), and other potential confounders. **Methods:** This case-series study benefits from an existing cohort of women ($N = 882$), ages 40 years and older, diagnosed with and treated for their first primary invasive breast cancer at the Geisinger Health System between 2001 and 2007. Data were obtained from Geisinger's electronic health record; the diagnosis atypical ductal hyperplasia (ADH) was the outcome of interest. Women were categorized according to their BMIs as normal, overweight or obese. All statistical tests were two-sided and analyses were performed using SAS version 9.3 (SAS Institute, Inc. Cary, NC). **Results:** The prevalence of biopsy prior to the diagnosis of breast cancer was not statistically different among the three groups of women. The risk of hyperplastic lesions was elevated for overweight ($OR = 13.58$; 95% $CI = 1.03-178.84$; $P = 0.047$) and obese ($OR = 14.76$; 95% $CI = 1.09-200.09$; $P = 0.043$) women relative to women in their ideal body weight. The association between BMI and other risk factors for breast cancer did not reach the level of statistical significance. **Conclusions:** Preliminary results suggest that excess body weight increases the risk of hyperplastic lesions in postmenopausal women. Biologically, this observed risk may be the result of higher serum levels of unbound estradiol in overweight and obese women. Future larger studies should validate our findings.

Keywords: Breast cancer, Obesity, Hyperplastic lesions

doi:10.3121/cmr.2010.943.ps3-12

PS3-13:

Patient Barriers to Mammography Identified During a Mammogram Reminder System

Adrienne Feldstein, MD, MS¹; Nancy Perrin, PhD¹; Ana Rosales, MS¹; Jennifer Schneider, MPH¹; Mary Rix, RN, BS, CCRP¹; Russell Glasgow, PhD²

¹Kaiser Permanente Northwest; ²Kaiser Permanente, Colorado

Background/Aims: Patient mammogram reminders are effective at raising screening rates, but patient screening barriers remain. We evaluated patient characteristics available in electronic medical records that might be associated with mammogram completion after a reminder program, and identified patient-reported barriers that could help to explain response differences. **Design:** Retrospective cohort study using electronic medical record data and a sub-group patient survey. Setting/Participants: Participants were female Kaiser Permanente Northwest HMO members aged 50-69 who were 20

months past their last mammogram (index date) and who were the target of a reminder intervention (Total $N = 4,708$). A mailed survey was completed by 336 of 665 (50.5%) members of the sub-group that received the survey. **Intervention:** “Mammogram due soon” post card 20 months after last mammogram, followed by a sequence of up to 2 automated phone calls and 1 live call for non-responders. Main outcome measure: Mammogram completion at 10 months following index date. **Results:** The characteristics associated with lower mammogram completion rates were age < 60 ($OR = 0.69$; $p < 0.0001$); health plan membership < 5 years ($OR = 0.81$; $p = 0.019$); family income $< \$40,000$ per year ($OR = 0.77$; $p = 0.018$); and obesity ($OR = 0.67$; $p < 0.0001$). Nearly 47% of the cohort was obese. Among survey respondents, being “too busy” ($OR = 0.54$; $p = 0.048$), feeling “embarrassed” ($OR = 0.09$; $p = .001$) and “too much pain” ($OR = 0.41$; $p < .001$) were associated with lower likelihood of mammogram completion. Obese women were more likely to report “too much pain” from mammograms compared to non-obese women (31.3% of 160 vs. 18.8% of 176; $p < 0.01$). Younger women were more likely to endorse that they believed in preventive screening (84.4% vs. 71.6%; $p < 0.05$), but also that they were “too busy” (19.1% vs. 6.4%; $p < 0.001$) and had more worries about mammogram accuracy (2.5 vs. 2.3 on 5 point scale; $p < 0.05$) than older women. Pain mediated the relationship between obesity and mammogram completion rates (indirect effect = -0.111 , 95% $CI = -0.229$ to -0.025 , $p = .008$). **Conclusions:** Important barriers to mammogram completion remain even with effective reminder systems. Obesity is an especially prominent barrier due to patients concerns about pain associated with mammography. Interventions to address/reduce pain may help lower this barrier.

Keywords: Mammogram screenings, Mammogram reminder system, Patient screening barriers

doi:10.3121/cmr.2010.943.ps3-13

PS3-14:

Primary Care Providers Reported Use of Pre-Screening Discussions for Prostate Cancer Screening: A Cross-Sectional Survey

Maggie Gunter, PhD¹; Ann Von Worley, BA, RN¹; Herbert Davis, PhD¹; Shelley Carter, MPH, MCRP, RN¹; Lawrence Scholl, MPH²; Jacqueline Miller, MD³

¹LCF Research; ²Macro International Inc.; ³Centers for Disease Control and Prevention

Background and Aims: There is no scientific consensus for routine prostate cancer screening for men of average risk less than 75 years of age using the prostate-specific antigen test (PSA). However, suggested practice is for physicians to discuss the known harms and potential benefits of PSA testing and digital rectal exam (DRE) with their patients. We conducted a survey of primary care providers (PCPs) to determine their practice styles including discussion regarding prostate cancer screening; we also conducted a patient survey to learn whether their PCPs discussed PSA testing and DRE during a health maintenance exam (HME). **Methods:** PCPs in the Lovelace Health System were surveyed about their prostate cancer screening practices for average risk men. Based on their responses, they were divided into two groups: 1) routine screeners (RS), those who routinely order PSA testing without discussing potential harms and benefits and 2) informed decision makers (IDM), those who routinely discuss the PSA test and its benefits and harms. After the HME, patients were surveyed as to whether or not their PCP discussed the PSA test and its harms and benefits during their visit. **Results:** Among the physicians, 35.9% (23/64) were RS and 64.1% (41/64) were IDM. Of the RS patients, 82.3% (51/62) responded that their physician did discuss the PSA test compared to 71.2% (37/52) of the IDM patients ($p = 0.10$). For RS, 58.8% (36/62) of the patients stated their provider initiated the discussion, whereas 56.8% (30/52) of the IDM initiated it. Provider recommendations to have the test were higher for the RS (80.3%) (41/51) compared to 53.1% (20/37) of the IDM ($p = .0032$). **Conclusions:** By patient report, a higher percentage of RS providers discussed the PSA test with patients. As would be expected, most of the RS recommended the PSA compared to only one-third of the IDM providers. Further studies are needed to determine what other factors of physician practice styles relate to discussing or not discussing prostate cancer screening with patients, and how