challenges in design and execution, particularly for Medicaid providers. In order to find an alternative that overcame the challenges of P4P, CareOregon—a non-profit health plan serving Medicaid and Medicare enrollees in Oregon—developed the Care Support and System Innovation (CSSI) Program. An evaluation was conducted to assess the overall impact of the CSSI program in promoting comprehensive improvements in quality of care among network providers and the extent to which it offers a viable alternative to P4P. **Methods:** Data were collected from the 48 medical practices that participated in the CSSI Program between 2005 and 2007. Data collection strategies included site visits, telephone interviews, document review and a web-based survey. Qualitative data were coded and analyzed using a grounded theory approach, emphasizing identification of themes judged meaningful to the study subjects. Quantitative data were analyzed through the use of descriptive statistics and chi-squared analysis. **Results:** The evaluation found that the CSSI Program successfully addressed many challenges inherent in P4P. The CSSI Program successfully engaged providers, fostered a culture of quality improvement within its provider network, and provided practices with the opportunity to develop sustainable and innovative solutions to address quality concerns. Success of the program was attributed to CareOregon's focus on building relationships while providing technical assistance and responsive funding. **Conclusions:** The CSSI Program has provided valuable support for improving quality of health care delivery and encouraged innovation among participating providers. The model provides a viable alternative for overcoming many of the challenges faced by pay-for-performance and other quality improvement programs.

Keywords: Quality of health care delivery, Care Support and System Innovation Program, Quality improvement assessments

doi:10.3121/cmr.2010.943.ps3-24

PS3-25:

Raising Literacy and Capacity for Incontinence and Skin Care in Dementia: Needs Assessment of Family/Friend Caregivers and Healthcare Providers

Jody Jackson, RN, BSN¹; Donna Bliss, PhD, RN²; Cheri Rolnick, PhD, MPH¹; Kenneth Hepburn, PhD³; Casey Arntson²

¹HealthPartners; ²University of Minnesota; ³Emory University

Background and Aims: Incontinence can occur in Alzheimer's disease or dementia (AD). Skin damage is frequently a physical complication. Few AD support programs directly address these problems. The purpose of this study is to assess and better meet the needs of informal caregivers and healthcare providers of persons with AD about issues of incontinence and perineal skin care. **Methods:** Audiotaped focus groups or interviews are being conducted with informal (family/friend) caregivers (goal n=56) and healthcare providers including physicians and nurse practitioners (NPs) (goal n=10) to determine needs for raising literacy and care capacity related to incontinence and skin damage. This assessment addresses health literacy content (e.g., knowledge, skills) as well as attitudes and preferred methods of receiving content. Results of this assessment, review of literature of current evidence and recommendations by expert clinical consultants will guide development of an informational and supportive intervention. **Results:** To date, 30 informal caregivers have participated in discussions (mean age 60; 75% female; 50% of care recipients are incontinent). A total of 11 healthcare providers participated in discussions (8 MDs, 2 NPs, 1 Pharmacist; mean age 53; 64% female; mean years of professional experience 23). Literacy needs of informal caregivers include prevalence and reasons why incontinence occurs in AD, when to discuss with healthcare provider and terms to use, types and photos of skin damage that can occur, indications for various absorbent and skin care products, and "step by step" instructions on ways to manage. Literacy needs of providers include evidenced-based recommendations for non-pharmacological management strategies, terms used by families and patients to refer to incontinence, referral or follow-up visit with specialist nurse, and utilization of provider (primary care, geriatrics) who has established relationship with patient. Preferred methods of learning/dissemination for both caregivers and providers include informational brochure, FAQ sheet, and guide to skin care products. Few preferred computer-based materials. **Conclusions:** Family/friend caregivers and

healthcare providers of individuals with AD have indicated health literacy needs related to incontinence and associated skin damage and have identified preferred ways of addressing these needs.

Keywords: Incontinence and perineal skin care, Health literacy, Caregivers and health providers

doi:10.3121/cmr.2010.943.ps3-25

PS3-26:

Reasons for Not Achieving Optimal Secondary Prevention Care for Coronary Artery Disease

Thomas Kottke, MD, MSPH¹; James Smith, MD²

¹HealthPartners Research Foundation; ²HealthPartners Medical Group

Background and Aims: Adherence to four secondary prevention goals (not using tobacco, LDL measured and <100 mg/dl within one year, blood pressure <140/90 at last visit, and daily aspirin) significantly improves outcomes for patients with coronary artery disease (ICD 9 410-414). However, a significant proportion of patients fail to achieve these four goals. We wished to determine the reasons why patients with CAD failed to achieve care goals in our multispecialty medical group. **Methods:** We abstracted 170 randomly-selected records of patients with CAD who failed to achieve care goals. **Results:** The two most frequent reasons for failure to achieve care goals were: the care team overlooked care needs at the time of an acute care visit (n=98), and the patient failed to return for follow-up (n=28). Other less frequent reasons were: The patient was offered treatment but declined (n=14); the clinician changed a medication in response to a value that was out of range, and the patient was not yet due for follow-up (n=10); and, the patient was not invited back for follow-up (n=9). The following reasons were observed in only one case each: Co-morbid conditions did not justify aggressive goals; the ordering physician failed to follow-up on a completed test; and, the patient was intolerant of statins. Eight patients failed to meet the optimal CAD care goal for other unique reasons. **Conclusion:** Providing clinical care teams with the tools to better track patient care needs (e.g., electronic patient identification and follow-up systems) could significantly reduce secondary prevention care gaps in our medical group.

Keywords: Coronary Artery Disease, Patient care outcome goals, Secondary prevention care

doi:10.3121/cmr.2010.943.ps3-26

PS3-28:

Telemedicine Specialty Consultation in a Medically Underserved Community

Ted Palen, PhD, MD, MSPH¹; Mike Bodily, MBA²

¹Colorado Permanente Medical Group; ²Kaiser Permanente Health Plan of Colorado

Background and Aims: During the current national health care reform discussions, health information technologies (HIT) are receiving a lot of attention as means of lowering health care costs, improving access to health care, reducing errors during the delivery of healthcare, and improving health outcomes. Patients in rural or medically underserved areas find it difficult to obtain and maintain adequate access to health care. Telemedicine (TM) services can exchange video and audio information between healthcare providers and patients in different locations. For individuals in rural communities TM may improve access to care through the use of novel care delivery models. We are conducting a proof of concept (POC) study to evaluate the use of TM consultation between a primary care clinic in a medically underserved region of southern Colorado and a specialty care (endocrinology) clinic in Denver. **Methods:** This is a prospective health services POC study. We are taking advantage of a "natural experiment," the expansion of primary care services into a medically underserved area of southern Colorado, to examine the requirements for establishing TM services between a rural primary care clinic and metropolitan based specialists. **Results:** Technical requirements included establishing network capabilities (512 kbps ISDN, H.320), 2 Mbps IP (H.323 or SIP), wireless LAN connection,