reported that hyperglycemia preferentially increases intracellular generation of angiotensin (Ang) II. However, circulating RAS is down regulated in diabetes implying the accelerated intracellular RAS mechanism as a major contributor to diabetic cardiomyopathy. In this study, we determined expression of RAS components and their effect on cardiac cells to give us a venue to intervene. **Methods:** Diabetes was induced in Sprague-Dawley rats and C57/BL6J mice with injection of streptozotocin for 5 days and verified by sustained blood glucose levels >15 mmol/L. Control mice received buffered saline alone. One week after diabetes induction the mice hearts were removed and perfused by the Langendorff method. Myocytes were isolated with enzymatic dispersion and centrifugation. RAS proteins were determined by real-time PCR and Western analysis. These included angiotensinogen, renin, angiotensin converting enzyme, AT1, AT2, and ACE 2. Angiotensin II was purified by reverse-phase chromatography and quantified by competitive ELISA. **Results:** Among cells obtained from diabetic hearts, expression of AGT (3.5+/- 0.8 fold), renin (2.4 +/- 0.4), and AT1 (2.6+/- 0.3) was significantly increased compared to cells from control hearts ($p < 0.05$, ANOVA). No significant change in the gene expression of ACE (1.2+/- 0.4), ACE2 (0.97+/- 0.2), and AT2 (1.2+/- 0.2), was observed. Increased expression at the protein level for AGT (2.2 +/- 0.1), renin (1.9 +/- 0.08), and AT1 (2.3+/- 0.2) was also observed by Western analysis. No significant changes in the protein levels of AT2, ACE, and ACE2 were observed. AngII levels in cardiac myocytes were determined by quantitative ELISA, which demonstrated significantly enhanced levels of AngII (140+/- 10 fmol/mg protein) synthesis in diabetic mice compared to controls (20 +/- 10 fmol/mg protein). **Conclusions:** This study suggests that local activation of the RAS in diabetes has a significant role in the development of cardiac dysfunction. Demonstration of renin expression in cardiac myocytes and enhanced expression in diabetic conditions represent a particularly significant finding. The latter suggests that locally produced, not circulating renin has a major role in cardiac Ang II generation and subsequent cardiomyopathy.

Keywords: Renin-angiotensin system, Diabetes, Cardiac dysfunction

doi:10.3121/cmr.2010.943.ps1-05

PS1-06:

The Impact of Treatment Intensification and Medication Adherence in Achieving Blood Pressure Control Among Coronary Artery Disease Patients

Colleen Ross, MS¹; Thomas Maddox, MD²; Ella Lyons, MS¹; Heather Tavel, BS¹; David Magid, MD, MPH¹

¹Kaiser Permanente Colorado; ²Denver VA Medical Center

Background and Aims: Among coronary artery disease (CAD) patients, control of hypertension reduces rates of myocardial infarction and mortality. Yet, BP control among CAD patients remains sub-optimal in clinical practice. Few prior studies have had the longitudinal data necessary to assess the role of treatment intensification and medication adherence in achieving optimal control. **Methods:** In this retrospective cohort study, we evaluated baseline and 1-year BP control rates among 15,102 HTN patients with an incident diagnosis of CAD between January 1, 2003 and December 31, 2006. Among those patients with sub-optimal baseline BP control, we assessed the contribution of treatment intensification and medication adherence to achieving 1-year BP control. Sub-optimal BP control was defined as systolic BP =140 mmHg and/or diastolic BP =90 mmHg (for those with diabetes or renal disease, sub-optimal control was defined as SBP =130mmHg and/or DBP =80mmHg). Treatment intensification was defined as an increase in BP medication dosage, the addition of another BP medication, and/or a switch of medication that resulted in a higher bioequivalent dose. Medication adherence was calculated using the proportion of days covered by medication prescription, and defined as a PDC > 80%. Logistic regression was used to determine the contribution of treatment intensification and medication adherence on achieving 1-year BP control, adjusting for baseline patient variables. **Results:** At the time of CAD diagnosis, 9,513 (65%) had sub-optimal BP control. At 1 year, 3,628(38%) of these subjects achieved BP control. Treatment intensification occurred more often among those patients who achieved 1-year BP control than those who did not (54% vs. 42%, p -value <0.0001). Medication adherence was similar between patients who

achieved 1-year BP control and those who did not (87% vs.86%, p -value 0.10) during the year post-CAD. In adjusted analyses, treatment intensification was significantly associated with 1 year BP control (OR 1.42 95%CI 1.39, 1.45). In contrast, medication adherence was not significantly associated with 1 year BP control (OR 1.02, 95%CI 0.87, 1.16). **Conclusions:** Treatment intensification, but not medication adherence, was significantly associated with 1-year BP control among a newly diagnosed CAD population. Interventions to improve treatment intensification may improve BP control and patient outcomes.

Keywords: Blood pressure, Treatment intensification, Medication adherence

doi:10.3121/cmr.2010.943.ps1-06

Diabetes and Obesity

C-C2-02:

The High Five for Kids Study: An Intervention to Improve Primary Care to Prevent Childhood Obesity

Elsie Taveras, MD¹; Steven Gortmaker, PhD²; Ken Kleinman, ScD¹; Kathy Mitchell, MD³; Sarah Price, MPH¹; Lisa Prosser, PhD⁴; Sheryl Rifas-Shiman, MPH¹; Matthew Gillman, MD¹

¹Harvard Pilgrim Health Care Institute; ²Harvard School of Public Health; ³Harvard Vanguard Medical Associates; ⁴University of Michigan Health System

Background and Aims: The preschool age group is ripe for clinically-based obesity prevention interventions. **Objective:** To assess the extent to which a primary care-based obesity prevention intervention (Ix), compared with the usual care control condition (UC), resulted in a smaller age-associated increase in body mass index (BMI) and improvement in specified obesity-related behaviors over the first year of a planned 2-year intervention. **Methods:** High Five for Kids is a cluster-randomized controlled trial in 10 primary care pediatric offices in Massachusetts. We randomly assigned 5 practices to the Ix condition and 5 to UC. Eligible children were age 2.0 - 6.9 years at baseline with BMI > 95th percentile or 85th-<95th percentile if at least one parent was overweight. The major components of the intervention were restructuring primary care based on the Chronic Care Model, stage-based motivational counseling by nurse practitioners, and stage-matched educational modules targeting TV, fast food, and sugar sweetened beverages. We enrolled 486 participants of whom 459 (94%) had primary outcomes measured at 1 year. We used intention-to-treat analysis, adjusted for baseline socio-demographic characteristics and corrected for clustering. **Results:** Mean (SD) age of participants was 4.9 (1.2) years; 29% had an annual household income < \$50K; 57% were white, 20% black, 16% Hispanic. Compared with UC, Ix participants had slightly smaller change in mean BMI from baseline to 1 year (-0.18 kg/m²; 95% CI: -0.44, 0.08). Ix participants decreased their TV viewing (-0.41 hrs/day; 95% CI: -0.68, -0.13) and fast food intake (-0.17 servings/week; 95% CI: -0.34, 0.00) more than UC, but there was no effect on sugar sweetened beverage intake (-0.20 servings/day; 95% CI: -0.51, 0.11). In post-hoc stratified analyses, the intervention had a greater effect on BMI change among participants living in households with incomes <\$50K (-0.79 kg/m²; 95% CI: -1.45, -0.13) than in higher income households (0.01 kg/m²; 95% CI: -0.34, 0.36). **Conclusions:** After a 1-year intervention period, High Five for Kids was effective in reducing TV viewing and fast food intake but did not significantly reduce BMI.

Keywords: Obesity prevention, Obesity-related behaviors, Obesity interventions

doi:10.3121/cmr.2010.943.c-c2-02