C-C3-05:

Characteristics of Patients Not Up-To-Date for Colorectal Cancer Screening (CRCS) and Who Decline Participation in a Trial to Increase Screening

Beverly Green, MD, MPH¹; Andy Bogart, MS¹; Jessica Chubak, PhD¹; Sally Vernon, PhD²; Ching-Yun Wang, PhD, MS³

¹Group Health Research Institute; ²University of Texas Health Science Center at Houston; ³Fred Hutchinson Cancer Center

Background/Aims: Screening decreases colorectal cancer incidence and mortality. However, almost half of eligible adults are not screened at recommended intervals. SOS: Systems of Support to Increase Colorectal Cancer Screening and Follow-up (R01CA121125) is a 4-arm randomized controlled trial that uses a systems-based approach to increase CRCS. We describe characteristics of patients who were not up-to-date compared to those current for CRCS and those who refused to participate in the study. **Methods:** The setting was 21 primary care clinics at Group Health, in the Pacific Northwest. Between 2008 and 2009, automated databases were used to identify patients age 50-73 with no life-threatening diseases. We used automated data to ascertain their prior CRCS status. Those not up-to-date were called to confirm their eligibility and willingness to participate in the trial. We collected automated and survey data to determine covariates related to not being up-to-date and refusing participation. **Results:** Based on automated data alone, 54.4% (28,508/52,556) of age-eligible patients were not up-to-date for CRCS. Those not up-to-date were younger, had fewer preventive care visits and screening tests for other cancers, lower levels of expected clinical need (a co-morbidity score), and less continuity of care with one physician. Patients not up-to-date for CRCS were randomly selected (n=15,451) to receive recruitment letters and telephone calls to confirm eligibility. Patients refusing to participate (N=6646) were younger, had lower levels of education, were more often from minority racial or ethnic groups, on Medicaid, and had fewer preventive care visits and other cancer screening tests than those who enrolled. They also more frequently reported they had never had any type of CRCS compared to participants (60.9% vs. 46%, $P<.001$). **Conclusion:** Similar to other studies, patients not up-to-date for CRCS are more often from underserved groups and get less preventive and cancer screening care. Unfortunately, these same factors are associated with refusal to participate in a trial to increase CRCS. Studies requiring verbal informed consent will have a hard time testing the effectiveness of interventions to increase CRCS for those most needing it.

Keywords: Colorectal cancer screening, Randomized controlled trial, Recruitment

doi:10.3121/cmr.2011.1020. c-c3-05

PS2-19:

Provider and Survivor Perspectives on Components, Timing and Source of a Colorectal-specific Cancer Survivor Care Plan

Jody Jackson, RN, BSN¹; Laurie Van Arman, LPN¹; Cheri Rolnick, PhD, MPH¹

¹HealthPartners

Objective: While much has been written about the value of cancer survivor care plans (SCPs), little has been done to examine what stakeholders want in such tools. A project is underway to assess essential elements of a colorectal-specific cancer SCP from the perspective of healthcare providers and survivors. **Methods:** To date, 42 healthcare providers (oncology physicians and nurses, social workers, surgeons, radiologists and primary care) have been invited to participate in a two-part online survey. In addition, 54 survivors (44 from HealthPartners and at least 10 from other Minnesota Cancer Alliance organizations) were surveyed via mail. Survey content included desired components (elements rated as very, somewhat, not at all important), and when and how to introduce SCPs. **Results:** Of the 42 providers surveyed, 66% (28 of 42) responded to Part One and 55% (23 of 42) to Part Two. To date, 29 survivors have responded. Over 90% (26 of 28) of providers rated the following content as very important: tumor site and

stage, long-term side effects, signs of recurrence, frequency of surveillance tests and oncology follow-up visits, as well as information on sexual issues and depression/anxiety. About 90% (26 of 29) of survivors categorized the following as very important: diagnostic tests received, surgical history, chemotherapy received, short-term and long-term side effects and how to manage them, and frequency of surveillance tests and oncology follow-up visits. Over half (14 of 20) of providers would introduce an SCP at treatment completion. About one-third (10 of 29) of survivors reported the tool should be introduced at time of diagnosis, almost one-third (9 of 29) at treatment completion and 21% (6 of 29) at beginning of treatment. The most frequently reported source of SCP information among both providers (83-91%, 19-21 of 23) and survivors (59-79%, 17-23 of 29) were oncology physicians or nurses. **Conclusions:** Providers and survivors have identified components essential to a colorectal cancer survivor care plan. Knowledge of the fundamental components from these key stakeholders will help focus care plan efforts to better meet the needs of survivors.

Keywords: Survivor care plans, Colorectal cancer

doi:10.3121/cmr.2011.1020.ps2-19

PS2-38:

Breast Cancer Survivorship Program for Rural Pennsylvania: Communicate, Connect, and Support

Rebecca Stametz, MPH¹; Azadeh Stark, PhD²; Adele Spegman, PhD, RN¹

¹Geisinger Medical Center; ²Geisinger Health System

Background: Advances in early detection and treatment of breast cancer has changed this disease from a fatal to chronic health condition. Recognition of informed decision-making and choices as the tenet of living well by the survivors motivates the development of credible and easily accessed information resources. Little is reported regarding the challenges specific to women in rural areas. Presently, available resources to encourage self-care are ineffective for women in our region, as evidenced by above the national average age- and stage-adjusted disease specific mortality rate. We have implemented a pilot project that leverages on technology to partner survivors and professionals that emphasize self regulation and health management. **Methods:** Development of this web-based interactive resource is directed by an advisory board. The content, which embodies the regional cultural norms and limitations of resources, addresses the immediate and the long-term personal, family and health needs of the survivors. Features include strategies and cues to guide self-management, problem solving and decision-making. Cancer survivorship plans are promoted, with suggestions regarding developing and implementing. The website is hosted by Geisinger and linked to the-Patient-Provider Communication portal for readily accessible networking among survivors, the oncology team, and primary care providers. Alpha and beta testing are planned; once vetted and published; tracking metrics will be evaluated for ongoing assessment. **Results:** An advisory board that constitutes of a balanced number of survivors and health professionals has been assembled. During the initial face-to-face meeting, members confirmed the intricacies of health and life needs of the survivors in our region. The board endorsed the value of a technology-leveraged interactive communication portal with providers. Furthermore, it was unanimously agreed to first focus on survivors who have received their initial treatment at Geisinger; however, the importance of a long-term objective to expand the network to survivors external to Geisinger and to survivors from all cancers clearly resonated. **Conclusion:** This innovation actualizes our commitment to patient-centered care by aligning convenience, technology, and personalized care. It serves as a template that readily can be modified with potential application to other geographic areas.

Keywords: Breast cancer, Survivorship, Website

doi:10.3121/cmr.2011.1020.ps2-38