adjusted for sex, age, BMI percentile, race/ethnicity and clinic department. **Results:** Overall, blood pressure was measured at >80% of children's visits to pediatric and family medicine clinics. The predicted probability of a blood pressure recorded at a primary care visit increased steadily with age (from .81 in 3-5 year olds to .89 in 15-17 year olds, p for trend <.0001). Of the 60% of visits where children had a recorded BMI, children with a BMI =95th percentile were slightly less likely to have a blood pressure recorded compared to BMI 85th <95th, and BMI<85th (91%, 93%, 94%, respectively, p <.0001). There were no differences by sex or race/ethnicity in the adjusted prevalence of documented blood pressure measurement. **Conclusions:** Blood pressure measurements are documented at a large proportion of children's visits to pediatric and family medicine clinics. Younger children were less likely to have blood pressure measurements recorded than adolescents. The relative lower likelihood of blood pressure measurement in obese children warrants further investigation given the higher prevalence of hypertension in these children.

Keywords: Pediatric, Hypertension, Obesity

doi:10.3121/cmr.2011.1020.c-b5-04

C-B5-03:

Linking Parent and Child Medical Records to Identify Youth at Risk for Familial Hypercholesterolemia

Catherine McNeal, MD, PhD¹; Debbie Godwin, BS¹; Linda Robinson, BS, MA¹; Dean Kjar, MS¹; Bhushan Mahajan, MS¹; Teresa Ponder¹; Anne Robertson¹

¹Scott & White Healthcare

Background: Despite consistent recommendations in numerous guidelines sponsored by professional societies, screening youth for familial hypercholesterolemia (FH) and/or those with a family history of premature cardiovascular disease (CVD) identifies only a small percentage of high-risk youth. By linking parent and child data in the virtual data warehouse, we can assess current screening practices, factors that impact screening practices and ultimately strategies aimed at improving childhood screening. **Aims:** the aim of this study is to compare cholesterol testing and screening rates in youth who have at least one parent covered under the same health plan with a history of CVD defined as a history of myocardial infarction, percutaneous coronary artery intervention (PCI) or coronary artery bypass surgery (CABG) and/or a parent with FH established by ICD-9 diagnoses or an abnormal cholesterol level. **Methods:** A cohort of youth who were 2-26 years of age anytime between service dates inclusive of 01/01/2001 and 12/31/2009 and who were covered as insured members under their parents' health plans within the Scott & White HMO were identified including a subset with any ICD-9 diagnosis code associated with hyperlipidemia. Parent-child linkages were created using a subscriber_id relationship code and relationship description. The parental cohort with CVD was identified from the CVRN Surveillance study as well as parents who have FH defined as an LDL-C of > 215 mg/dl or a total cholesterol > 300 mg/dl. **Results:** We found that adherence to guidelines for cholesterol screening of youth at a higher risk for premature CHD is no greater than screening rates for children with a lower risk who do not have an affected parent. Moreover, a family history of CHD and hypercholesterolemia is rarely documented in a dependent's medical record. **Conclusions:** Cost-effective approaches to screening for FH in the pediatric population rely on a parental history of CVD and/or FH. However, this approach does not identify the majority of youth at risk for premature CVD. Universal screening of youth may be warranted.

Keywords: Hypercholesterolemia, Children, Screening

doi:10.3121/cmr.2011.1020.c-b5-03

C-B5-01:

Childhood Obesity: Risk Factors, Associated Morbidity and Service Utilization

Revital Gross, PhD¹; Daphna Doron, MD, MHA²; Yael Ashkenazi, MPH¹; Beatriz Hemo, MA²; Orit Ben Shoham, MHA²; Israella Nahshon, MHA²

¹Myers-JDC-Brookdale Institute; ²Maccabi Healthcare Services

Aims: To identify parent-level and child-level risk factors for childhood obesity, to study health consequences and medical service utilization among the overweight children. **Methods:** The study included families with children aged 4-11 years old, registered with Maccabi Healthcare Services at 2 community clinics in the Greater Jerusalem Area. During 2008, 409 mothers completed questionnaires and interviewers weighed and measured 714 children. There was a 50% response rate. Information on associated morbidity and utilization of Health Services was obtained from Maccabi's computerized database. **Results:** Approximately 30% (122 of 409) of the families had at least one overweight or obese child (BMI>85th percentile / BMI>95th percentile, respectively). Twenty percent of the children sampled were overweight or obese (overweight: 11.9%, 85 of 714. obese: 8.3%, 59 of 714). The variables with an independent effect on having at least one overweight child in the family were: father of Asian-African origin; immigration to Israel since 1995; non-academic education; a maternal age less than 40; a mother who had a history of being overweight and who reported lack of control over her eating habits. Variables with an independent effect on children's obesity/overweight were: age below 7 years old; eats a broad variety of foods (not fussy); has an 'inactive' lifestyle; eats the main daily meal at a daycare center or afternoon enrichment center at least once a week. Among the overweight children, mainly in the 7-11 age groups, a higher rate of associated morbidity (particularly asthma, orthopedic problems and dermatological conditions) was documented. The service utilization rates were also higher among overweight children compared to children within the normal weight range. **Conclusions:** Pro-active medical monitoring of overweight or obese children must be encouraged even if they do not present other symptoms. Parents and caregivers should be provided with information regarding the risk factors for obesity, as some of those are modifiable. It is recommended that a follow-up study be conducted with a national representative sample of children to verify the findings.

Keywords: Obesity, Child, Risk factors

doi:10.3121/cmr.2011.1020.c-b5-01

C-B5-02:

A Parent Failure Modes and Effect Analysis: Parent Teams Can Identify Problems and Propose Interventions

Joann Wagner, MSW¹; Kathleen Walsh, MD, MSc²; Juliet Moretti, BA¹; Terry Field, DSc¹; Kathleen Mazor, Ed, DMS¹; Jennifer Donovan, PharmD³; Abir Kanaan, PharmD³

¹Meyers Primary Care Institute-UMass Medical School, Fallon Clinic, FCHP; ²UMass Medical School, Meyers Primary Care Institute; ³MCPHS, Meyers Primary Care Institute

Background: Parents of many children with cancer administer highly complex chemotherapy regimens at home which are error prone. One way to identify and repair potential error prone processes is with a Failure Mode and Effect Analysis (FMEA). An FMEA is a systematic, team-based approach to understanding the ways a process can fail and developing interventions; it has been used in hospitals for decades but has never, to our knowledge, been used with a team of parents. **Methods:** The objective of this study was to identify aspects of the home medication use processes at risk for error and propose interventions using an FMEA with a group of parents. We recruited parents of children with cancer taking home medications. FMEA steps included: selection of a high-risk process; diagram the process; brainstorm potential failure modes; rate and prioritize failure modes; identify root causes; and redesign the process. The target high-risk process was parent/caregiver administration of 6-mercaptopurine following a change in medication dose. Parents were trained in FMEA methodology using materials at a 6th grade reading level and a real-life example. Failure modes were prioritized using ratings for severity, frequency of occurrence, and detectability. **Results:** The process diagram developed by the parent-team included 12 steps. The highest priority failure modes were: (1) shared instructions by the parent who attended the clinic visit regarding change in dose are misunderstood by the other caregiver(s); (2) administration of chemotherapy without realizing that another caregiver already administered it; (3) parent misunderstanding of physician's instructions about the change in dose; and (4) the parent who attended the clinic visit shares incorrect instructions regarding the change in dose with other caregivers. During the