Background/Aims: Inadequate laboratory monitoring of high-risk medications is a medical error associated with preventable adverse drug events. Patient failure to complete ordered laboratory tests is not uncommon. We conducted in-depth interviews with patients who did not complete a recently ordered laboratory test to identify barriers to test completion. **Methods:** The target population was patients in a large multispecialty group practice who failed to complete an ordered laboratory test. We conducted semi-structured interviews with 18 patients who did not complete a monitoring test order related to 1 of 7 high-risk medications (angiotensin converting enzyme inhibitors (ACE) inhibitors, angiotensin II receptor blockers (ARBs), digoxin, phenytoin, statins, thyroid replacement hormone, or valproic acid) between July 2008 and October 2010. For comparison, we also interviewed 5 patients who completed all ordered laboratory tests. The interviews occurred in person when possible (n=17) as well as by phone (n=6). All interviews were recorded and transcribed and analyzed using a grounded theory approach. **Results:** The laboratory monitoring discussions included the following themes: physician communication about lab tests, patient comprehension of lab test purpose, patient attitude toward lab tests, and barriers to test completion. The most prominent barrier to test completion was forgetting. Patients often did not realize they had missed an ordered test. When patients did acknowledge missing an ordered test, they often noted that they did not encounter any barriers besides forgetting. Regarding potential interventions, patients reported that understanding the indication for the tests would not help reduce non-completion. Several patients understood the reason for the test, but still failed to complete it. While some thought it was important to understand the reason for the test, others trusted their physician and did not believe it was essential to understand. Some patients were using an online personal health portal system offered through the practice that included information about upcoming tests. **Conclusions:** Forgetting was the most prominent theme for non-completion of ordered test. Patients felt that providing timely (near desired test completion date) patient reminders, either by phone or by e-mail, would likely improve completion rates of laboratory monitoring for high-risk medications, while providing more information about the test indication would not.

Keywords: Laboratory monitoring, Qualitative research, High-risk medication

doi:10.3121/cmr.2011.1020.ps2-13

C-B3-01:

Building Toward a Patient-Centered Culture: Insights and Tactics from Literature, Health System Stakeholders, and Patients

Sarah Greene, MPH¹; Leah Tuzzio, MPH¹; Dan Cherkin, PhD¹

¹Group Health Research Institute

Background: The degree to which medical care is patient-centered is frequently cited as a key attribute of healthcare quality. However, the precise definitions, skills and behaviors that underlie patient-centered care remain somewhat elusive. **Methods:** We used a pragmatic literature review, focus groups with patients, and interviews with key delivery system and health plan leaders as well as primary care teams to guide development of an operational definition of patient-centered care. We then sought specific opportunities to partner with the health system leadership and disseminate this work, in order to mobilize our entire organization to find ways to become more patient-centered. As part of the focus groups and interviews, we gathered feedback about a proposed definition of patient-centered care that could be adopted organization wide. **Results:** The literature review identified key attributes of patient-centered care at the system level, clinical level and interpersonal level. Both focus groups and interviews showed concordance in their views of what patient-centered care was--how it was defined, and how it was manifested by our doctors and our entire organization. We have utilized both evidence from the scientific literature and input from our members and practitioners to identify specific strategies and intervention opportunities for our health system. The field of patient-centered care research is fertile, however a clear taxonomy is needed to describe how patient-centeredness relates to related constructs such as activation, empowerment, doctor-patient communication, trust, and shared decision-making. Promising interventions exist, but few have been replicated.

Conclusions: Systems, and individuals within systems have opportunities to further develop and exemplify a patient-centered culture. Identifying specific attributes of delivery system design, communication, clinician training and optimal care transitions are among the areas that are ripe for improvement. Key issues that need to be addressed for successful cultural and system change and the highest priorities for research to support such change will be highlighted.

Keywords: Patient-centered care, Quality improvement, Culture

doi:10.3121/cmr.2011.1020.c-b3-01

PS2-21:

Explanation to Understanding: Practices of the Highest Performing Physicians by Patient Survey

Tom Janisse, MD, MBA; Kaiser Permanente

Background: The MD-Patient Communication Study aims were the best practice communication behaviors of physicians during outpatient clinic visits; and to collect physician perspectives of communication behaviors. Researchers have consistently found the top predictors of overall patient satisfaction are quality of the physician-patient relationship and contributing communications. There is limited understanding however, of the range of specific behaviors in the interaction associated with positive and negative patient perceptions and reactions. **Methods:** In phase 1, we conducted a naturalistic, observational study of fifty-five, Permanente Primary Care physician-patient visits using videotape recordings, and incorporating patient and physician reactions to the tape. The physicians, who practiced in Los Angeles and Honolulu, spanned the three strata of high, medium, and low historical patient satisfaction scores. In phase 2, a standardized six-question set was posed, in semi-structured, 60-minute interviews, to 77 of the highest-performing physicians on this patient survey, including: 20 of the highest performers (top 5%) from the LA and Honolulu groups; and 42 and 15, respectively, of the highest performing physicians in Portland and Oakland. These interviews were audio taped with permission, transcribed and coded for patterns. **Results:** Explanation to achieve understanding was the most commonly cited activity in answer to a 'physicians' role in their patients' healing,' and in what constitutes a "therapeutic moment in the visit." Pattern dimensions include: context – trust, relationship; communication – connection, dialogue; content – visual explanation; role – patient ownership; and support – reassurance, encouragement. Representative verbatims: "It's how the information is interpreted and packaged, and what's emphasized, and how it's customized to the individual;" "I spend a fair amount of time explaining physiology in their terms so they understand why we're doing what we're doing, or what's happening and what to expect;" "If you don't have a relationship with your patient, then you don't have a really effective treatment protocol." **Conclusions:** With national health care reform's new Prevention Codes calling for "Discussion," and patients' dependence on explanation to understand and follow through with treatment recommendations, describing how high performing physicians create high patient satisfaction, is of great importance.

Keywords: Explanation, Understanding, Relationship

doi:10.3121/cmr.2011.1020.ps2-21

Diabetes, Obesity, and Physical Activity

PS2-27:

Near-Normalization of Glucose and Microvascular Diabetes Complications: Data from ACCORD and ADVANCE

Patrick O'Connor, MD, MPH¹; Faramarz Ismail-Beigi, MD, PhD²

¹HealthPartners Research Foundation; ²Case Western Reserve School of Medicine

Purpose: To compare results of clinical trials that assessed the impact of near-normalization of glucose on microvascular complications in type 2 diabetes. **Methods:** ACCORD (N=10,234) and ADVANCE (N=11,140) tested the hypothesis that near-normalization of glucose reduces microvascular complications in adults with established type 2 diabetes. Difference in