obese had an OR of 1.40 (95%-CI 1.35-1.45; p for trend<0.001) for any injury or pain in the lower limbs compared to children and adolescents with normal body weight after adjustments for sex, age, race/ethnicity, and socioeconomic factors. Similarly, extremely obese children and adolescents had increased relative odds for: fractures (OR 1.26, 95%-CI 1.20-1.33); sprains/strains (OR 1.47, 95%-CI 1.36-1.59), dislocations (OR 1.41, 95%-CI 1.36-1.47), and pain (OR 1.45, 95%-CI 1.24-1.70). For all injuries except dislocations, the odds ratios increased with increasing degree of overweight/obesity (p for trend <0.001), with underweight children/adolescents having lower relative odds of all injuries and pain compared to the normal weight group. **Conclusions:** Overweight, obese, and extremely obese children/adolescents are more likely to experience lower extremity injuries and/or pain than are their underweight and normal weight peers. While the precise mechanism underlying these associations remains unclear, lower extremity injuries can reduce the physical activity in weight groups that need to increase activity levels to reduce or manage their weight.

Keywords: Obesity, Childhood, Injuries

doi:10.3121/cmr.2011.1020.c-c2-02

Plenary III-01:

HMORN Research on Pediatric Hypertension and Obesity: Predictors, Care, and Costs: Design of a New Pediatric Cohort

Kenneth Adams, PhD¹; Emily Parker, PhD, MPH¹; Nicole Schneider, BA¹; Heather Tavel, BS²; Malini Chandra, MS²; Mary Becker, BA¹; Maureen Peterson, BA¹; Karen Margolis, MD, MPH¹; Nancy Sherwood, PhD¹; Elyse Kharbanda, MD, MPH¹; Alan Sinaiko, MD³; David Magid, MD, MPH²; Matthew Daley, MD, MPH²; Joan Lo, MD²; Patrick O'Connor, MD, MPH¹

¹Health Partners Research Foundation; ²Kaiser Permanente; ³University of Minnesota Medical

Background: The increasing numbers of children or adolescents having elevated blood pressure (BP) is an important public health problem, but little information is available regarding patterns of care or resource use implications in this population. We established a data cohort of children and adolescents to study these relationships. **Methods:** We have implemented a longitudinal cohort study based on routinely collected electronic medical record data from 3 large health plans. The study will ultimately extend from calendar year 2007 through 2012. We have currently extracted subject data through 2009. We will continue to accrue new subjects through 2012 and to add data for existing subjects in future years. The source population consists of children and adolescents (3-17 yr) enrolled in the 3 health plans. Beginning in Jan 2007, age-eligible subjects enter the cohort at the time of the first clinic visit in which an eligible BP is measured (i.e., collected during an outpatient visit and associated with a corresponding height measure). Subjects are being followed forward in time with longitudinal data collection. Demographic, clinical, and administrative data include age, gender, race/ethnicity, BP, height, weight, diagnoses, medications, and cost of care. We have conducted preliminary analyses to demonstrate the accrual of subjects into the cohort over years 2007 through 2009, and describing the gender, age, and racial composition of the study population. **Results:** As of the end of 2009, the cohort included 330,880 children and adolescents. At entry, 58.1% were 3-11 years of age, whereas 41.9% were 12-17 years. Subjects were 50.2% female. Race was 41.0% white (combining Hispanics and non-Hispanics), 8.1% African-American, 12.8% Asian/Pacific Islander, 16.3% other, and 21.8% missing. **Conclusions:** We have created a large pediatric cohort of subjects using data collected from health plan electronic medical and administrative records at several HMORN sites. These de-identified data include longitudinal BP, BMI, and utilization data suitable to test hypotheses related to correlation of BP and BMI over time, the influence of BMI on BP, patterns of care provided to those with elevated BP and BMI, and the influence of elevated BP and BMI on resource use.

Keywords: Predictors, Electronic medical records, Pediatric hypertension

doi:10.3121/cmr.2011.1020.plenary-iii-01

C-C2-04:

Diabetic Patients and Risk – It's Not About the Numbers

Barry Saver, MD, MPH¹; Kathleen Mazor, EdD¹; Lee Hargraves, PhD²

¹UMass Medical School/Meyers Primary Care Institute; ²UMass Medical School

Background: The major, common adverse health outcomes for patients with diabetes come from macrovascular, not microvascular, disease, including death, heart attack and stroke. Personalized risk information might help patients make better decisions about prioritizing health behavior change. In this study, we explored diabetic subjects' beliefs about diabetes-related risks and sources of information along with reactions to presentation of personalized risk information. **Methods:** We recruited 56 English- and Spanish-speaking adults with diabetes and at least one other major cardiovascular risk factor from a community health center and an academic family medicine practice. In in-depth interviews, we explored sources of risk information, risk-reduction activities, and behavioral intentions. Subjects also ranked 6 diabetes-associated adverse outcomes based on perceived risk, reviewed personalized risk predictions from the UKPDS Outcomes Model, and re-ranked these perceived risk of outcomes. We explored their reasons for changing/not changing risk rankings. Qualitative analysis was used to develop themes and concepts underlying subjects' risk perceptions and reactions to risk information. **Results:** Virtually all subjects believed at least some diabetes-related adverse outcomes were modifiable or preventable. A substantial minority expressed a fatalistic view that at least some were not preventable, while others felt that risk factor control meant an outcome was "not going to happen." A very common theme was that a "warning shot" would occur before many outcomes, providing time and impetus to change bad habits. Provider urging was the other commonly-cited motivator and about 20% of subjects cited each as motivators for past change. While providers were cited as a key source of information, vicarious experience with relatives and acquaintances was cited just as often and 3/4 of subjects felt they knew themselves and their risks better than providers or risk models. Subjects reported the personalized risk information was interesting, but <1/3 changed their rankings of mortality (nearly always initially ranked least likely) to match the model predictions (nearly always the highest probability). **Conclusions:** Our subjects often based their risk perceptions on anecdote and gut feelings, not medical information. Personalized risk information does not appear promising for motivating behavior change or even altering risk perceptions.

Keywords: Diabetes, Risk perception

doi:10.3121/cmr.2011.1020.c-c2-04

C-B5-05:

Early Childhood Obesity Prevention in Primary Care: Opt-In Versus Opt-Out Recruitment Strategies

Margaret Rukstalis, MD¹; Chelsie Hauer, BA¹

¹Geisinger Health System

Background: Although early childhood obesity prevention programs are needed in primary care to address the growing pediatric obesity epidemic, recruitment remains particularly challenging. Aims To describe opt-in versus opt-out recruitment strategies to invite parents of overweight/obese 4-8 year olds to participate in a family lifestyle modification obesity prevention study in primary care. **Methods:** Following IRB approval, traditional recruitment strategies were employed including flyers, doctor referrals, and public ads to invite parents of 4-8 year olds with a Body Mass Index (BMI) > 85th percentile for age and sex to call (Opt-IN) about participating in a 5-month primary care obesity prevention intervention. Electronic Health Records were searched and physician letters were sent to parent/guardian of eligible children who resided at zip codes within 1-hour commute of treatment site to invite them to call about the study. Opt In letters were mailed over 13-months for Cohorts 1-3 recruitment. For Cohort 4, identical physician letters were mailed except parents were asked to call if they were NOT interested in information about the study (Opt-OUT). After 10 days, study staff called to

explain the study and schedule an in person screening. **Results:** In 2008-2009, over 15,000 children ages 4-8 had at least one height and one weight recorded in the EHR at the same visit with BMI > 85th % for age, sex. 11,276 physician letters were mailed: 9,954 opt in letters over 13-months (Cohorts 1-3) and 1,322 opt out letters in 2-months (Cohort 4). Of the 170 families screened, 58/107 (54%) identified opt in letters as recruitment source; 49/63 (78%) families in Cohort 4 reported opt out letters introduced them to the study. Fifty-six families gave written informed consent and/or assent and were randomized to one of two groups – Group Family Behavior Modification (FBM) or Enhanced Information Mailing (EIM) group. Three FBM and two EIM families dropped before treatment started. **Conclusions:** Physician letters to parents of eligible children identified in the EHR were the most successful recruitment strategies for the primary care obesity prevention study. Opt-out physician letters shortened recruitment time and doubled the number of participants recruited for the final cohort.

Keywords: Obesity, Prevention, Childhood

doi:10.3121/cmr.2011.1020.c-b5-05

C-C2-05:

Primary Care Web-Based Lifestyle Intervention for Type 2 Diabetes: Randomized Controlled Trial to Improve Knowledge and Self-Care

Margaret Rukstalis, MD¹; Mary Ann Blosky, MHA, MS, RN¹; Howard Steinberg, MBA²; Mary Anglade, MBA²; Tammy Anderer, PhD, MSN, CRNP¹; Frederick Bloom Jr., MD, MMM¹

¹Geisinger Health System; ²Life Med Media

Background: Rapidly growing numbers of Type 2 Diabetes (T2D) in the US continues to escalate need for evidence-based primary care interventions to reduce complications and costs. Limited time, information, and lack of revenue for chronic disease management create gaps between national guidelines and primary care for diabetes. **Methods:** Of the over 20,000 diabetics were identified in Geisinger EPIC® EHR, 3166 T2D with diagnosis on problem list or ICD-9 code had HgA1c > 8.0%. Physician letters or emails were sent to invite them to call to Opt OUT if they did not want more information. After a 10 day period, staff called 1932 eligible T2D to invite them to a shared medical primary care visit to participate in a randomized controlled trial comparing a tailored 5-month web-based lifestyle intervention (dLifeG.com) to usual care. 166 Type 2 diabetics gave written informed consent and were randomized 1:2 to control group or intervention group. In <20 minutes with simple computer instructions, T2Ds in intervention group created password protected personalized website to set goals, view weekly lessons, and take interactive quizzes to improve diabetes knowledge and self-management. Weekly emails with lesson topics and links to the dLifeG.com were sent to intervention group participants. At end of study, control group will also have access to website intervention. **Results:** At midpoint, ~3/4th of the 100 T2 diabetics in intervention group were engaged [defined by the number of site page viewed (0-3090 pages), emails opened (0-100%), and quiz pages consumed (0-578)]. Mid and end of study changes in diabetes knowledge, HgA1c, Blood Pressure, and weight will be discussed. **Conclusions:** Web-based lifestyle interventions can be employed in primary care to engage a majority of diabetics with HgA1c > 8.0 in a self-management lifestyle modification intervention. Importantly, the study also helped identify diabetics who may need additional resources and assistance with chronic disease management.

Keywords: Diabetes, Intervention, Self-care

doi:10.3121/cmr.2011.1020.c-c2-05

PS2-01:

Factors Influencing Completion in the IDEA Diabetes Education Study

Kenneth Adams, PhD¹; Sarah Beaton, PhD²; Emily Parker, MPH, PhD¹; Jodi Lavin-Tompkins, RN, CNP¹; JoAnn Sperl-Hillen, MD¹; Ann Hanson, BS¹; Herbert Davis, PhD²; Omar Fernandes, MPH¹; Ann Von Worley, RN, BSHS²; Patricia Glasrud, MS, RDABQ¹; William Parsons, MS²; Vic Spain, PhD, DVM³

¹HealthPartners; ²LCF Research; ³Merck and Co, Inc.

Background/Aims: Participant non-adherence to study protocols can adversely affect clinical trials by reducing statistical power. Similarly non-adherence is an important issue in clinical practice. Ceasing to participate is one form of non-adherence. Subject characteristics appear to influence adherence. We evaluated relationships between subject characteristics and completion of educational sessions in IDEA, a clinical trial testing 2 modes of diabetes education in patients with sub-optimal diabetes control. **Methods:** IDEA eligibility criteria included Type 2 diabetes, A1c ≥ 7%, and no recent diabetes education. Subjects (n=623) were randomly allocated to one of 3 diabetes education treatment arms; individual education, group education, or a comparison group with no active intervention. Individual education consisted of three 1-hour sessions, following an AADE program. Group education was delivered in four 2-hour group sessions that emphasized patient interaction. Subjects were considered adherent if they completed all assigned sessions. At the end of the study, tracking data indicated completion rates of 72.0% (175/243) in the group education intervention and 86.1% (211/245) in the individual treatment intervention. We sought to identify demographic, psychosocial and clinical characteristics that might explain non-completion. We hypothesized that, within each arm, baseline health status (SF-12 mental composite and physical composite scores), depression (PHQ9), personality type (TIPI Big 5; extraversion, agreeableness, emotional stability, conscientiousness, and openness), and Hb A1c level were associated with completion. **Results:** In the group education arm, subjects scoring higher on the emotional stability scale were more likely to complete (p<0.05). Generally, demographic factors were more strongly associated with completion of group education than were psychosocial or clinical factors: women and older subjects were more likely to complete than men and younger subjects (gender, p=0.008; age, p<0.0005). Within the individual education arm, completion was predicted by higher physical health score (SF12-PCS; p=0.005), higher mental health scales (SF12-MCS, p=0.008), lower depression score (PHQ9, p=0.002), and lower Hb A1c (p<0.05). However, neither gender nor age was associated with completion. **Conclusions:** Factors related to study completion appear to differ between the 2 diabetes educational interventions. These results may be relevant to diabetes educational programs considering strategies to improve “no show” and lack of completion rates.

Keywords: Diabetes education, Group education, Adherence

doi:10.3121/cmr.2011.1020.ps2-01

PS2-06:

Regional Variation in Response to Diabetes Education in the IDEA Study

Sarah Beaton, PhD¹; JoAnn Sperl-Hillen, MD²; Omar Fernandes, MPH²; Ann Von Worley, RN, BSHS¹; William Parsons, MS¹; Herbert Davis, PhD¹; Ann Hanson, BS²; Emily Parker, MPH, PhD²; C. Vic Spain, DVM, PhD³

¹LCF Research; ²HealthPartners Research Foundation; ³Merck and Co., Inc.

Background/Aims: ‘Journey for Control of Diabetes: The IDEA Study’ is an ongoing randomized trial evaluating the efficacy of group diabetes education (GE) using US Diabetes Conversation Maps™ and conventional individual diabetes education (IE) among adults with sub-optimally controlled Type 2 diabetes (A1c ≥ 7%). The study is being conducted at 2 sites, Minneapolis, MN, and Albuquerque, NM. Because the 2 sites are regions with diverse demographic characteristics, the goal of this paper was to identify differences in the main outcome (A1c) between sites. **Methods:** In addition to A1c ≥ 7, study inclusion criteria were patients who were diagnosed with diabetes, and were diabetes education-naïve within the last one to two years. Consenting patients were randomly assigned to receive either: GE, IE or usual care (UC) as recommended by the primary care provider. All education was provided through each care system’s ADA-recognized education program. Follow-up A1c levels were obtained from electronic laboratory data. Differences in A1c levels were evaluated using Generalized Linear Models (SAS 9.2). **Results:** There were 623 patients enrolled: 337 from MN and 286 from NM. Sample demographics were: mean age of 60.0 (MN) and 64.0 (NM); percent female, 44.8 (151/337) for MN and 54.9 (157/286) for NM; race/ethnicity, 81.1% (276/337) white, 8.1% (27/337) black, 3.6% (12/337) Hispanic, and 7.2% (24/337) other for MN, and NM 45.6% (130/286) white, 2.5% (7/286) black,