Stevens-Johnson syndrome, toxic epidermal necrolysis, and angioedema, and extreme weight gain (EWG) among adults on atypical antipsychotics. **Methods:** Project work was divided among three teams. HealthPartners and Marshfield Clinic led the DILI team, Kaiser Permanente Hawaii and Kaiser Permanente Georgia led the SSR team and Group Health and Geisinger guided the EWG study. Teams met routinely via biweekly conference calls to coordinate their study efforts; monthly conference calls were conducted to coordinate the overall project, mark progress and discuss challenges and solutions. For each study, standardized case identification criteria were developed with input from iSAEC, expert panels, and medical literature. Potential cases were identified through electronic data searches using diagnoses, medication histories, laboratory test results, and other clinical data elements indicative of DILI and SSR during 2000-2009, or EWG from 2004-2009. Potential cases of DILI and SSR were abstracted to assess provisional case status and determine implicated drugs. Weight trajectories of suspected EWG cases were visually inspected to confirm EWG during atypical antipsychotic treatment period. **Results:** A total of 99 provisional cases of DILI, 41 cases of SJS/TEN and 56 provisional cases of angioedema were identified from the electronic records of the participating HMOs. For EWG, 249 cases were confirmed and an additional 341 were categorized as "possible" cases, with additional chart review and patient interview required for confirmation. Confirmation rates varied from 17% for DILI to 2-79% for SSR and 23-30% for EWG. **Conclusions:** The SAEC and HMORN study sites identified well-phenotyped cases with the targeted drug-induced SAEs of interest. The relative success of these efforts has been critical in the planning of a larger second study that will include analyses of genetic factors associated with provisional case SAEs.

Keywords: Drug-induced liver injury, Serious skin rashes, Extreme weight gain

doi:10.3121/cmr.2011.1020.ps1-18

PS1-23:

The Incidence and Epidemiology of Inpatient and Outpatient *Clostridium difficile* Infections at Kaiser Permanente

Jennifer Kuntz, PhD¹; Eric Johnson, PhD¹; Marsha Raebel, PharmD²; Amanda Petrik, MS¹; Xiuhai Yang, MS¹; Karen Glenn, BS²; Micah Thorp, DO, MPH¹; Nancy Neil, PhD, ICON; David Smith, PhD¹

¹Kaiser Permanente Northwest; ²Kaiser Permanente Colorado

Background/Aims: *Clostridium difficile* is the most common cause of healthcare-associated infectious diarrhea in the United States. However, few population-based epidemiologic studies of *C. difficile* infection (CDI) exist. Our aims were to: 1) describe the epidemiology of CDI among HMO members from two geographical regions; 2) extrapolate the incidence of CDI from these health plans to the U.S. population; and, 3) identify patient characteristics that predict inpatient versus outpatient identification of CDIs. **Methods:** We conducted a population-based, dynamic, retrospective cohort study among Kaiser Permanente Colorado (KPCO) and Northwest (KPNW) enrollees between June 1, 2005 and September 30, 2008. We identified incident CDIs and categorized infections based on identification in the inpatient or outpatient setting. We calculated incidence rates and extrapolated our estimates to the 2007 U.S. population. KPNW and KPCO regional electronic databases provided data on membership, pharmacy dispensings, demographics, clinical measures, and healthcare utilization. Logistic regression determined how strongly patient characteristics predicted inpatient versus outpatient identification of CDIs. **Results:** We identified 2,879 incident CDIs; 55% were identified in the outpatient setting. In 2007, our total population incidence rate was 13.5 CDI cases per 10,000 person-years; incidence increased with age. Extrapolated to the U.S. white-only population, we estimate that 220,000 cases of CDI occurred among persons 20 years or older in 2007. Baseline characteristics and healthcare utilization prior to identification of CDI in both settings were similar, although individuals with CDI identified in the inpatient setting were older and had greater comorbidity than individuals with outpatient-identified CDI. We identified few strong and independent predictors of setting, although risk for CDI identification in the inpatient setting was associated with an estimated glomerular filtration rate <15 (OR: 4.96; 95% CI: 2.73, 9.02), inflammatory bowel disease (OR: 2.41;

95% CI: 1.45, 4.02), an outpatient antimicrobial dispensing in the previous 180 days (OR: 2.67; 95% CI: 2.13, 3.35), and malignancy (OR: 2.08; 95% CI: 1.60, 2.72). **Conclusion:** The incidence of CDI among this population is substantial (13.5/10,000 person-years). Because the incidence of and risk for CDIs increases with age, the U.S. burden of CDIs will continue to rise as the proportion of older individuals in the U.S. population rises.

Keywords: *Clostridium difficile*, Incidence, Epidemiology

doi:10.3121/cmr.2011.1020.ps1-23

PS1-11:

Feasibility of Prospective Surveillance for Adverse Drug Events – Lessons Learned from a 12-Month Pilot Study

Sascha Dublin, MD, PhD¹; Taliser Avery, BS²; Craig Cheetham, PharmD, MS³; Robert Davis, MD, MPH⁴; Martin Kulldorff, PhD²; Lingling Li, MD²; Liyan Liu, MD⁵; Jeffrey Brown, PhD²

¹Group Health; ²Harvard Medical School and Harvard Pilgrim Health Care Institute; ³Kaiser Permanente Southern California; ⁴Kaiser Permanente Southeast; ⁵Kaiser Permanente Northern California

Background/Aims: When new medications are introduced, safety data are often sparse. Health plan data may be useful for active surveillance for adverse events. We conducted a 12-month pilot study to demonstrate the feasibility of prospective active surveillance using data from several large health insurers. This presentation will summarize logistical challenges and lessons learned. **Methods:** We studied the safety of generic divalproex sodium, an antiepileptic drug introduced in July 2008, compared to the branded drug. We performed monthly surveillance using the maximized sequential probability ratio test, comparing observed outcomes in new users of the generic drug to expected counts based on new users of the branded drug from January 2002 - June 2008. Programmers at four HMO Research Network sites extracted data monthly, ran quality control (QC) programs, and returned summary tables to the lead site. Medication exposure was identified from outpatient pharmacy dispensings. Outcomes identified from health plan utilization data included potential adverse effects (e.g. pancreatitis and liver disease) and efficacy outcomes (e.g. hospital or emergency department visits for seizures.) **Results:** We will discuss lessons learned related to data sharing and HIPAA compliance, timing and synchronization of data pulls, and QC activities. Obtaining IRB approval took 3.5 months, and protocol development took 7 months. Developing the analytic program required two rounds of corrections, one leading to meaningful changes in results. QC programs identified substantial differences in monthly extracts across sites. Some QC flags required only understanding differences across sites, but others revealed data problems requiring resolution at the site level. Overall, 14% (4/28) of monthly extracts across the four sites could not be used due to underlying data problems or lack of analyst availability for timely extracts. **Conclusions:** Prospective drug safety surveillance is feasible, but future projects should anticipate the need for longer timelines and multiple revisions to data pulls and analytic programs. They should develop approaches to handling missing data, because some data pulls will not be usable. QC activities should continue throughout study implementation. A knowledgeable, engaged lead at each site is vital to ensure accuracy and appropriate interpretation of each site's data.

Keywords: Antiepileptic medications, Drug safety, Prospective surveillance
doi:10.3121/cmr.2011.1020.ps1-11

C-C5-05:

Genetic Risk Factors: Idiopathic Thrombocytopenic Purpura following the First Dose of Measles-Mumps-Rubella Vaccination in Children

Mia Hemmes, MPH, CHES¹; Robert Davis, MD, MPH¹; Thomas Buttolph, MD²; Yandong Qiang, MD, PhD²; Deborah Nickerson, PhD³; Jill Johnsen, MD³; Christopher Carlson, PhD⁴

¹Kaiser Permanente Georgia; ²FDA; ³University of Washington; ⁴Fred Hutchinson Cancer Research Center